Supporting Information 2

Iodinated Meroditerpenes from a Red Alga Callophycus sp.


Serge Lavoie,^{†,‡} David Brumley,[†] Troy S. Alexander,[†] Christine Jasmin,^{†,‡} Franz A. Carranza,[‡] Kate Nelson,[§] Cassandra L. Ouave,^{§,¶,◊} and Julia Kubanek^{†,‡,*}


†School of Chemistry and Biochemistry and ‡School of Biological Sciences, Aquatic Chemical Ecology Center, Institute for Bioengineering and Biosciences, Georgia Institute of Technology, Atlanta, Georgia 30332.


§Department of Dermatology, ¶Center for the Study of Human Health, and [◊]Antibiotic Resistance


Center, Emory University, Atlanta, Georgia 30322


*julia.kubanek@biosci.gatech.edu
















Microdilution checkerboard analysis showing the combined effect of compounds **1–5** with oxacillin against CA-MRSA USA300, where the extent of inhibition is shown as a heat plot
















Microdilution checkerboard analysis showing the combined effect of compounds **1–5** with vancomycin against VISA, where the extent of inhibition is shown as a heat plot











Microdilution checkerboard analysis showing the combined effect of compounds **1–5** with vancomycin against VREF, where the extent of inhibition is shown as a heat plot