

The 2023 wildfire season in Québec: an overview of extreme conditions, impacts, lessons learned, and considerations for the future

Yan Boulanger^a, Dominique Arseneault^b, Annie Claude Bélisle^c, Yves Bergeron^{d,e}, Jonathan Boucher^a, Yan Boucher^f, Victor Danneyrolles^g, Sandy Erni^h, Philippe Gachon^h, Martin P. Girardin^a, Eliane Grantⁱ, Pierre Grondin^l, Jean-Pierre Jetté^o, Guillemette Labadie^k, Mathieu Leblond^l, Alain Leduc^{d,e}, Jesus Pascual Puigdevall^a, Martin-Hugues St-Laurent^b, Junior A. Tremblay^{m,n}, and Kaysandra Waldron^{b,a}

^aCentre de Foresterie des Laurentides, Canadian Forest Service, Natural Resources Canada, 1055 rue du Peps, Québec City, QC G1V4C7, Canada; ^bDépartement de biologie, chimie et géographie, Université du Québec à Rimouski, 300 allée des Ursulines, Rimouski, QC G5L 3A1, Canada; ^cConseil de la Première Nation Abitibiwinni, 45, rue Migwan, Pikogan, QC J9T 3A3, Canada; ^dDépartement des sciences biologiques, Université du Québec à Montréal, Case postale 8888, Succursale Centre-Ville, Montréal, (Québec) H3C 3P8, Canada; ^eUniversité du Québec en Abitibi-Témiscamingue 445 Boul de l'Université Rouyn-Noranda, QC J9X 5E4, Canada; ^fCentre de recherche sur la boréale, Département des sciences fondamentales, Université du Québec à Chicoutimi, 555, boulevard de l'Université, Chicoutimi, QC G7H 2B1, Canada; ^gGreat Lakes Forestry Centre, Canadian Forest Service, Natural Resources Canada, 1219 Queen Street East Sault Ste. Marie, ON P6A 2E5, Canada; ^hÉtude et Simulation du Climat à l'Échelle Régionale (ESCER), Université du Québec à Montréal, Case postale 8888, Succursale Centre-Ville, Montréal, QC H3C 3P8, Canada; ⁱCree Nation Government, 32 Amisk Street, Mistissini, QC G0W 1C0, Canada; ^jMinistère des Forêts, de la Faune et des Parcs, 5700, 4e avenue ouest, Québec, QC G1H 6R1, Canada; ^kDépartement des sciences biologiques, Université du Québec à Montréal, Case postale 8888, Succursale Centre-Ville, Montréal, QC H3C 3P8, Canada; ^lLandscape Science and Technology Division, Environment and Climate Change Canada, 1125 Colonel By Drive, Ottawa, ON K1S 5B6, Canada; ^mUniversité Laval, Pavillon Abitibi-Price 2405, rue de la Terrasse, Québec, QC G1V 0A6, Canada; ⁿWildlife Research Division, Environment and Climate Change Canada, 801-1550, avenue d'Estimauville, Québec, QC G1J 0C3, Canada; ^oArmagh, Québec, Canada G0R 1A0

Corresponding author: Yan Boulanger (email: yan.boulanger@nrcan.gc.ca)

Abstract

The 2023 wildfire season in Québec set records due to extreme warm and dry conditions, burning 4.5 million hectares and indicating persistent and escalating impacts associated with climate change. This study reviews the unusual weather conditions that led to the fires, discussing their extensive impacts on the forest sector, fire management, boreal caribou habitats, and particularly the profound effects on First Nation communities. The wildfires led to significant declines in forest productivity and timber supply, overwhelming fire management resources, and necessitating widespread evacuations. First Nation territories were dramatically altered, facing severe air quality issues and disruptions. While caribou impacts were modest across the province, the broader ecological, economical, and social repercussions were considerable. To mitigate future extreme wildfire seasons, the study suggests changes in forest management practices to increase forest resilience and resistance, adapting industrial structures to changes in wood type harvested, and enhancing fire suppression and risk management strategies. It calls for a comprehensive, unified approach to risk management that incorporates the lessons learned from the 2023 fire season and accounts for ongoing climate change. The study underscores the urgent need for detailed planning and proactive measures to reduce the growing risks and impacts of wildfires in a changing climate.

Key words: boreal forest, climate change, fire management, forest management, risk management

1. Introduction

Fueled by record-breaking warm and dry conditions (Barnes et al. 2023; ESCER 2023), the 2023 fire season that occurred in Québec (Canada) was one of extremes. By the end of October, more than 4.5 Mha of forest had burned throughout the province (SOPFEU 2023a), doubling the previous record set in 1989 (2.3 Mha). The total area burned within the com-

mercial forest, which also corresponds to the Intensive Protection Zone under fire management, reached approximately 1.1 Mha, the highest since 1923. These numbers are roughly equal to the area that has burned in the Intensive Protection Zone over the past 20 years combined. Extreme conditions for fire spread fueled the largest wildfire events ever observed in both the Intensive Protection Zone (460 kha) and

the northern protection zone where fire management is extensive (1 Mha) (CWFIS 2023). On 1 June, due to the numerous and fast spreading wildfires, the provincial authorities responsible for fire management in Québec declared that they had reached preparedness level 5 (CIFFC 2023), meaning that the situation had created extreme demands for provincial firefighting resources. Such a preparedness level resulted in the mobilization of national and international resources to assist the *Société de protection des forêts contre le feu* (SOPFEU), the agency responsible for fire management and suppression in Quebec (CIFFC 2023). International help was particularly timely as the whole country was at National Preparation Level 5 from 11 May to 7 September 2023, making interprovincial resource sharing very difficult. The situation was particularly hazardous as dangerous fires were burning near communities, leading to a significant number of fire-related evacuations, most notably by First Nation communities (Canadian Forest Service 2023). Additionally, the air quality was severely compromised, threatening the health of a large proportion of the population of the province, living up to several hundreds of kilometers away from the blazes (CBS News 2023). The dense smoke plumes originating from the 2023 fires in Québec also led to air quality alerts on multiple occasions for large areas in adjacent provinces as well as in northern United States (The New York Times 2023). Eventually, the smoke plumes would cross the Atlantic and cause hazy skies in western Europe (Leibniz Institute for Tropospheric Research 2023).

The 2023 fire season in Québec took many by surprise, as the previous decade had been particularly quiet in terms of wildfires (CIFFC 2023). However, several studies had warned of increased fire activity as a result of increased anthropogenic climate forcing, with potential dire consequences, notably for Québec's forest sector (Bergeron et al. 2010; Gauthier et al. 2015; Chaste et al. 2019) as well as on infrastructures and communities (Erni et al. 2021). Although consequences of these fires were immediate for communities, public safety, and industries, such an extensive and record-breaking fire season will no doubt have longer-term impacts on several important aspects of Québec's forest ecosystems, economy, and society. In the following sections, we report on the extent to which this season was exceptional in the context of historical variability. Furthermore, we summarize the impacts the 2023 wildfire season had and will have on (i) the forest sector, (ii) fire management, communities, and infrastructures, (iii) wildlife and their habitats, and (iv) First Nations in the province. We also provide potential avenues to mitigate these impacts.

2. The 2023 fire season in Québec: a timeline and a historical perspective

2.1. The 2023 weather conditions that fueled this extreme fire season

The province of Québec, with over 4.5 million hectares burned in 2023, was the most affected in terms of total area burned in Canada. Most fires occurred in the western and northwestern parts of the province, specifically within the

Eastern James Bay and Eastern Subarctic Homogeneous Fire Regime (HFR) zones, with 12% (the highest annual rate of any HFR in Canada since at least 1980) and 2.5% of their areas burned, respectively (Fig. 1).

Québec experienced its warmest May to August period since 1950, with maximum daily temperatures reaching record highs (Fig. 2). Although overall precipitation was near normal, the Eastern and Western James Bay HFR zones had one of their driest seasons on record, with the areas along the James Bay area experiencing Canada's driest anomalies (Suppl. Mat. S1.1). June saw the highest precipitation deficits, especially in the northwestern and northern parts of the province, persisting for most of the summer along the James Bay coast (Suppl. Mat. S1.2). Temperatures from April to July were consistently above average across the province, peaking in June with areas in the northern half of the province experiencing anomalies of over +3 °C, up to +5 °C (Suppl. Mat. S1.2). An early snowpack melt in April (Suppl. Mat. S1.3), leading to the lowest May snow water equivalent since 1950 (Fig. 2), facilitated the early onset and intensity of the wildfire season. Fire-weather index (FWI) for May through August reached levels significantly above the 1991–2020 normal for the province (Fig. 2). The Eastern James Bay and Eastern Subarctic and Western James Bay HFR zones saw the highest FWI anomalies, especially in June and July (Suppl. Mat. S1.2), the period during which most fires burned.

Unusual conditions of atmospheric blocking conditions have occurred during the spring of 2023 over the North Atlantic, corresponding to the maximum values of seasonal blocked days over the 1950–2023 period. When a blocking occurs a combination of negative/positive precipitation/temperature anomalies might lead to higher chances of getting forest fires and drought conditions, specially during spring and fall seasons (Zhao and Liu 2019; Wazneh et al. 2021; Antokhina et al. 2023). Overall, these conditions fueled an intense wildfire season that lasted from late May through September, peaking during three intervals (1–12 June, 19–28 June, and 3–15 July) where daily burned areas often exceeded 100 000 ha, coinciding with sustained high to extreme FWI values (Fig. 2). Of note, early June thunderstorms triggered over 120 wildfires, rapidly spreading due to high FWI values (Fig. 3). Persistent dry conditions from June to mid-July (Fig. 2) supported ongoing and new fires in the Eastern and Western James Bay and Eastern Subarctic HFR zones. Fire progression decreased after mid-July as FWI values dropped in the Eastern James Bay HFR zone (Fig. 3), where most active fires were located.

2.2. The 2023 fire season relative to the historical range of variability

The comparison of weather conditions and fire activity in 2023 with those of the past is essential to better understand their exceptionality in the context of climate change. An analysis by the World Weather Attribution initiative (Barnes et al. 2023) concluded that both the intensity (FWI values averaged over a 7-day period) and the severity (cumulative daily severity rating; Van Wagner 1987) of weather conditions from the beginning of the fire season through the end of July 2023

Fig. 1. Location of the 2023 wildfires in Québec (14 April to 1 October) as mapped by the *Direction de la protection des forêts* of Québec's *Ministère des Ressources Naturelles et des Forêts*. Only fires above 1000 ha are mapped. The Intensive fire protection zone is located south of the commercial forest northern allocation limit. Transects sampled for fire history in the James Bay area (see [Section 2.2](#)) are also shown. This figure was created using ArcGIS Pro v3.1.2 and assembled from the following data sources: Homogeneous Fire Regime (HFR) zones were retrieved from [Boulangier et al. \(2014\)](#). High-voltage powerline data are retrieved from [Ministère des Ressources naturelles du Québec \(2019\)](#). Basemap is from World topographic Canadian Style (<https://www.arcgis.com/sharing/rest/content/items/6d0ed88458c6429d99331260fb7bf2b0/resources/styles/root.json>). First Nation location was retrieved from Canadian Government (<https://open.canada.ca/data/en/dataset/b6567c5c-8339-4055-99fa-63f92114d9e4>). Public forests were retrieved from <https://www.donneesquebec.ca/recherche/dataset/unite-d-amenagement>. Fire location for 2023 was retrieved from <https://www.donneesquebec.ca/recherche/dataset/feux-de-foret/resource/b222e70c-26db-4af2-b24e-16b87e4acab7>. The commercial forest northern allocation limit was retrieved from <https://www.donneesquebec.ca/recherche/dataset/limite-territoriale-des-forets-attribuables>.




Fig. 2. Anomalies (expressed as standard deviations) for various meteorological parameters for the May–August 2023 period compared with the 1991–2020 normals for each Homogeneous Fire Regime zone (see Fig. 1), as well as for all zones combined (Québec). Ranking of 2023 values (from 1: highest to 74: lowest) against the 1950–2023 period is also shown for each parameter. FWI: Fire-weather index; SWE: snow water equivalent; Tmax: maximum daily temperature; Tmean: mean daily temperature; Tmin: minimum daily temperature; TotPrcp: total precipitation. Data were calculated from ERA5, the fifth generation ECMWF reanalysis (Hersbach et al. 2020).




Fig. 3. Daily area burned (in thousands of ha, gray bars: less than 30 kha burned; orange bars: 30–50 kha burned; red bars: 50–100 kha burned; dark red bars: >100 kha burned) within each homogeneous fire regime overlapping Québec during the 2023 fire season. Burned areas are estimated through M3 hot spots (CFS 2023). Daily fire-weather indices spatially averaged for the whole HFR zone and as assessed from ERA5 reanalyses (Hersbach et al. 2020, black lines) are also shown for the same period.





Fig. 4. Drought Code (DC) severity trends in Québec's boreal forests for early May, mid June, and early August (1900–2023). Map resolution: 0.5 degrees. (a) Ranking of DC severity for 1 May 2023, against the historical DC severity for the same date over a 124-year period. Regions in red represent instances where the daily DC level in 2023 closely approached the historical severity level at each grid point (refer to the legend on the right), while white dots denote record-high levels for 2023. (b–c) Similar to (a) but focusing on 15 June and 1 August. (d–e) Middle-row plots: time series illustrating the daily DC average for June encompassing the Eastern James Bay and Eastern Subarctic zones. The gray shading represents 90% confidence bands accounting for spatial autocorrelation. Horizontal dashed lines mark the severity of 2023. Bottom-row plots: analyses of extreme DC severity trends, estimated as the rate of occurrence (per year) of extreme drought years using a kernel approach (bandwidth parameter $h = 15$, based on Mudelsee 2002; Mudelsee et al. 2004). The gray shading represents 90% confidence bands for risk estimates. Additional details regarding the methodology are available in the Supplementary Information. Maps were created using ArcGIS v10.5.1 for Windows and ESRI spatial data (ESRI 2017).



were respectively at least twice and seven times more likely because of ongoing anthropogenic climate change. These results are in line with several other analyses that have concluded that many wildfire-weather metrics are being modified by climate change in Québec, including lengthier fire seasons (Jain et al. 2017), more severe and frequent extreme fire-weather indices (Jain et al. 2022), and drier fuels (Ellis et al. 2022). Most of these changes are actually occurring in northwestern and northern Québec, the very region where the 2023 fire activity was concentrated.

We used the Drought Code (DC), as measured from the FWI system (Van Wagner 1987) to assess trends in long term drought conditions in the province. DC records of 2023 showed that, as of 1 May in northern Québec, this fire season exhibited the driest conditions in 124 years (Fig. 4) and showed unprecedented drought severity across the province by mid-June (Vincent et al. 2020; ESCER 2023). The Eastern Subarctic and Eastern James Bay HFR zones experienced June

DC values that surpassed or equaled historical records, indicating severe drought. Previous research (Girardin et al. 2009) highlighted a mid-20th century decrease in extreme droughts in Canadian forests due to increased precipitation, but an increase in drought occurrences in northern forests due to warming. Since 2010, the Eastern Subarctic zone has seen more extreme drought events than in the 1970–1980s, aligning with earlier periods (1920–1960). The 2023 events may indicate an ongoing rise in drought severity in northern regions, with a pause in the 1970s. In the Eastern James Bay HFR zone, a post-1980s upward drought trend suggests a potential reversal of the declining trends observed up to 2002 (Girardin et al. 2009; Girardin and Wotton 2009). In this area, significant positive trends in summer DC values between 1950 and 2020 were observed, meaning an increase in drought conditions over southern shorelines of the Hudson Bay, from Manitoba towards Québec HFRs (see ESCER 2023). Similar trends are also observed in British Columbia (Parisien et al. 2023).

Over the past several centuries, variations in drought severity and frequency have significantly influenced fire regimes in Québec. Research shows that fire activity is closely linked to temperature, precipitation, vegetation, and human activities, highlighting the complex relationship between climate change and fires (Carcaillet et al. 2001; Remy et al. 2017; Girardin et al. 2019). To understand the 2023 wildfires, it is essential to differentiate between the wildfire histories of the Intensive Protection Zone and Northern Protection Zone, as historical fire drivers differed between these zones.

In the Intensive Protection Zone, burn rates were high during the Little Ice Age (around 1250–1850 AD; Gennaretti et al. 2014) and the early 20th century (Drobyshev et al. 2017; Danneyrolles et al. 2021; Chavardès et al. 2022). However, they remained low from 1940 until the extreme 2023 season, coinciding with the growth of the modern forest industry (Boucher et al. 2017a; Tymstra et al. 2020). The 2023 fire season was the most active in public commercial forests since 1923 (Fig. 5a), but when compared to long-term records, it still falls within the natural range of variability (Chavardès et al. 2022). About 7.4% of the western part of the Intensive Protection Zone burned in 2023, contributing to a decade-long mean annual burn rate of ~0.7%, which is still within the historical range of 0.6%–1.3% from 1750 to 1950 (Chavardès et al. 2022).

Conversely, in the Northern Protection Zone, particularly its western portion, fire activity in 2023 differs significantly from its recent range of variability. This region is among the most pyrogenic in Québec and the circumpolar boreal zone. Employing a methodology by Héon et al. (2014), fire size was reconstructed using dendrochronology along transects (Fig. 1; Erni et al. 2017; Shakeri 2024). The fire regime within the Northern Protection Zone has been consistently active since 1800, marked by regularly occurring extreme fire years without a long-term trend (Fig. 5b; Erni et al. 2017; Shakeri 2024). Significant fire years in this region include 1847 (during which 90 km of the transects burned), 1882 (104 km), 1906 (104 km), 1922 (140 km), 1941 (96 km), 1989 (136 km), and 2013 (102 km). However, 2023 surpassed all these with 208 km burned transects, making it the most substantial fire year in the last 224 years.

Overall, these analyses confirm that the 2023 fire season is markedly distinct from those of the last century, in terms of severe weather conditions and areas burned, and that these conditions are getting more frequent in Québec. The fire activity, fueled by these conditions, is unprecedented in at least 220 years of record in the Northern Protection Zone. Despite being within the long-term natural range of variability within the commercial forests, the extent of these fires ranks among the highest of what has been recorded in the past century. Further global warming could exacerbate the ongoing trends identified above and ultimately lead to a 3- to 5-fold increase in annual area burned by the end of the 21st century in the province (Boulanger et al. 2014). Such increases in fire activity would strongly modify the boreal forest ecosystem (Chaste et al. 2019; Boulanger and Pascual 2021; Boulanger et al. 2022a), affect its ability to conduct sustainable forest management (Gauthier et al. 2015; Pau et al. 2023), substantially increase the province's fire management and suppression costs (Hope

et al. 2016), increase infrastructure and communities' exposure to short fire interval (Erni et al. 2021; Arseneault et al. 2023), and modify species-at-risk habitats (Tremblay et al. 2018; Leblond et al. 2022; St-Laurent et al. 2022). Impacts generated by the 2023 fire season and those projected in the upcoming decades because of climate change are presented in the following sections.

3. Impacts of the 2023 fire season


3.1. Effects on the forest sector

Québec's forest sector, a key part of the province's socio-ecosystem, faces significant challenges due to the 2023 wildfires, as well as the expected increase in such events in the coming decades. In 2020, this sector generated \$CAN 20.6 billion in revenue and contributed \$5.1 billion to the real GDP (NRCan 2020). Despite a decline over the past two decades (NRCan 2023), the sector remains crucial for the stability of many small, remote, and mono-industrial communities in Québec. The 2023 fire season induced important losses in silvicultural investments and will strongly impact timber supply over the coming decades. Drastic reductions in annual allowable cut (AAC) will occur in management units that were most hardly hit by the 2023 fires (Forestier en Chef 2023), which will span over several decades (see also Suppl. Mat. S3).

Furthermore, a conservative estimate of at least 300 000 ha of commercial forests might suffer from regeneration failures (Le Devoir 2023a; see Suppl. Mat. S4) because these stands were immature (<60 years) with an insufficient regeneration potential (Splawinski et al. 2019). Unless these areas are planted, they will likely remain unproductive for decades. These young stands are primarily associated with the cumulative historical impacts of harvesting and wildfires over the last 50 years (see Suppl. Mat. S4—Table S4.2). Considering the current capacity of the forest sector to establish plantations (~50 kha per year), bringing back these areas into production could take several years, a huge budget (probably several billions of dollars) and imply technical challenges at a level never seen before (e.g., building of new forest roads, shortages in seedlings, nursery, and planting labor). Economic consequences of these impacts reside in the substantial losses in silvicultural investments, including many plantations. We estimate that ~80 000 ha of plantations have burned in 2023 (Table S4.2).

To alleviate fire-induced losses, burnt forests with mature trees can be targeted for salvage logging. However, considering the extent of 2023 fires, and that mature forests represent a fraction of what has burned, only ~10%–20% of burned stands are likely to be salvaged in the upcoming year. As such, salvage logging will not offset the long-term deficit in timber supply induced by the 2023 fires. Additionally, recently burned trees sought by salvage logging are rapidly affected by degrading agents (e.g., wood-boring insects and wood checking) that reduce the economic value of postfire timber, and thus the profitability of such practices (Saint-Germain and Greene 2009; Boucher et al. 2020). Salvage logging, following a wildfire, acts as a second disturbance in a brief period. This combination of fire and logging can signif-

Fig. 5. (a) Annual area burned (AAB) and number of fires within the Intensive Protection Zone (IPZ) in Québec between 1923 and 2023 (retrieved from the archives of the *Ministère des Ressources Naturelles et des Forêts du Québec*). (b) Distance burned along the Billy Diamond and Trans-Taiga roads in northern Québec (as shown in Fig. 1; total of 640 km sampled). The years indicated correspond to fire events covering 90 km or more.



icantly alter the recovery and provision of various ecosystem services (Lindenmayer and Noss 2006; Leverkus et al. 2020). Salvage logging affects numerous aspects like tree seedbeds, seedling density, woody debris, biological legacies, water conditions, and soil properties (Leverkus et al. 2018, 2020). These influences have short- to long-term effects on the forest's structure, composition, diversity, and dynamics (Purdon et al. 2004; Nappi and Drapeau 2011; Thorn et al. 2018).

The immediate and extended impacts of the 2023 fires on Québec's forest sector reflect the numerous warnings issued by the scientific community in the past two decades. It was shown that climate-induced increases in wildfire activity posed a serious threat to achieving sustainable forest management goals (Gauthier et al. 2015; Forestier en Chef 2021), with wood shortages and regeneration failures representing a primary climate-induced risk for Québec's forests. With the expected surges in wildfires, Québec's commercial forests will face an increasing proportion of immature forest at the expense of harvestable forest stands, despite a pos-

sible increase in productivity in the northern forest (Pau et al. 2022; Wang et al. 2022; Danneyrolles et al. 2023). In addition, simulation experiments suggest that enhancing vegetation productivity may result in a positive feedback loop with increased fire activity, ultimately yielding little net benefit over the long term (Chaste et al. 2019). A business-as-usual strategy could imply permanently lowered timber supply (by more than 60% in some regions) to avoid fire-induced timber shortages (Forestier en Chef 2021). An increase in fire activity is also likely to amplify broadleaved tree regrowth at the expense of the preferred conifers (Boulanger and Pascual 2021), increasing salvage logging (Forestier en Chef 2021) and management costs (Cyr et al. 2022). This scenario could lead to a "manager's dilemma", where the significant social costs of intensively managing forests, such as replanting conifers in areas with failed postfire regeneration, must be balanced against the increased vulnerability of these forests to future wildfires. Also, frequent softwood timber supply shortages, higher stumpage prices, and decreased export value of forest-

related products could seriously affect the province's economy and potentially prompt the devitalization of forest communities in the long term (Williamson et al. 2007).

3.2. Impacts on the fire management agency, communities, and infrastructure

With an overwhelming number of simultaneous and intense wildfires, the 2023 fire season really tested Québec's fire management agency's (SOPFEU) operational limits, which are known to cap out at about 30–40 active fires a day. Indeed, in the Intensive Protection Zone only (Fig. 1; also refer to Cardil et al. 2019), the number of active fires jumped from 21 to 132 on 1 June, marking the start of 45 consecutive days with over 30 active fires. These numbers greatly exceed the historical (1994–2022) average of 14 ± 17 (mean \pm SD) days with over 30 active fires per year. As a result, the exceeding fires burned freely until reinforcements from the Canadian Armed Forces, and other provincial and international fire management agencies started to arrive on 5 June, adding 50 people to the 434 local resources already at work. External reinforcements peaked on 28 June with 994 people, while local resources peaked at 643 on 12 June. The lack of readily available resources at the provincial and national levels to respond to the workload of early June, coupled with the inefficient or unsafe conditions of fire suppression activities due to the extreme fire weather (Hirsch and Martell 1996), likely contributed to rapid spread of these fires, sometimes threatening communities and infrastructures. This led SOPFEU to prioritize fires threatening human lives and (or) infrastructure deemed essential to public security, that were numerous, over those simply threatening the forest, including silvicultural investments and standing timber (Cardil et al. 2019).

As climate change intensifies, scenarios like the 2023 fire season, where fire management capacities are overwhelmed, are expected to become more frequent. This is due to projected increases in fire occurrences, area burned, conditions favorable to fire spread, and days with intense fires that impede suppression efforts (Wotton et al. 2017). Recent work indicated that this would have a direct impact on both wildland firefighters and airtankers workload (i.e., number of hours worked doing fire suppression activities) (Boulanger et al. 2022b; Boucher et al. in preparation). The 2023 fire season is an example of these impacts, with a total firefighters (local and external) workload that summed up to 755 648 h, representing more than six times the historical (1994–2018) average of $117\,017 \pm 121\,530$ (mean \pm SD) h, and more than 1.5 times the maximum of 477 024 h observed in 2010 (Boucher et al. in preparation; 2023 data provided by SOPFEU). Airtankers flew a total of 3219 h fighting fires in 2023, more than three times the historical (1994–2018) average of 981 ± 740 (mean \pm SD) h, and just over the observed maximum of 2005 with 3096 h (Boucher et al. in preparation; 2023 data provided by SOPFEU). The 2023 fire season observed workload for both firefighters and airtankers fall within the range of expected future workload for the end of this century (2071–2100) under Representative Concentration Pathway (RCP) 8.5, under which $474\,300 \pm 211\,680$ (mean \pm SD) h for firefighters and 5832 ± 4212 (mean \pm SD) h for airtankers are pro-

jected (Boucher et al. in preparation). As extreme fire-prone conditions are expected to become more frequent, the 2023 wildfire season is thus a reminder that costs associated with fire management and suppression will greatly increase in the upcoming decades (Hope et al. 2016), thus putting more communities and infrastructures at risk.

Many communities faced direct fire threats and had to swiftly react to fires that were rapidly progressing towards critical infrastructures during the 2023 fire season. As a last barrier of defense, approximately 45–69 km (or 226 ha) of fire breaks (where vegetation was stripped to the mineral soil at a width of approximately 50 m) were created in a hurry to directly protect seven communities (Chapais, St-Lambert, Normétal, Chibougamau, Oujé-Bougoumou, Mistissini, and Lebel-sur-Quévillon, data provided by the *Direction de la Protection des Forêts* of the *Ministère des Ressources Naturelles et des Forêts du Québec*). These conditions also led to an unprecedented number of wildfire-related evacuations, with over 38 700 people evacuated, including many from First Nation communities (CFS 2023). Large northern Québec communities like Sept-Îles (population of about 25 000) and Chibougamau-Mistissini (population of about 10 000) were among those evacuated. Some towns, such as Lebel-sur-Quévillon, experienced multiple evacuations, adding stress to the residents, local public safety agencies, and officials. On 9 June, the province announced a financial help of \$1500 to each evacuated household (Gouvernement du Québec 2023a). Based on the number of evacuees per communities and the number of people per household (Statistics Canada 2023a), we estimated that about 14 509 residences were evacuated, for a total toll of this financial help estimated to 21.8 mol/L\$. The mental health toll was evident, as seen in the resignation of the mayor of Chapais in November 2023 due to post-traumatic stress disorder caused by the intense wildfire crisis management (La Presse 2023). Remarkably, there were no casualties directly linked to the fires. However, there were significant losses in terms of infrastructure, including forest machinery (Le Soleil 2023) and First Nation' infrastructures on traditional territories (Grant 2023). Indeed, approximately 1154 structures (min = 1, max = 627 structures per fire; Microsoft Canadian Building Footprint (2020) and Gouvernement du Québec (2023b)), mainly cabins, were in the path of 64 wildfires this season, leading to the destruction of many (J. Boucher, personal observations).

The significant wildfire activity in the James Bay region (Fig. 5b) consistently poses threats to communities and crucial infrastructures in Northern Québec, particularly the La Grande hydroelectric complex, a vital part of the province's energy network (Erni et al. 2017; Arseneault et al. 2023). This complex, contributing 40% of Québec's electrical power, comprises 11 hydroelectric stations and related infrastructures like high-voltage power lines, roads, airports, and residences. In 2023, many fires impacted strategic areas of Hydro-Québec, crossing major power lines that serve the province's most populated regions (Fig. 1), resulting in multiple shutdowns of high-voltage lines (Le Devoir 2023b). A similar situation occurred in 2013, leading to power outages affecting major areas, including Montreal, Québec's largest city, disrupting the subway system and affecting hundreds of thousands

of people. With anticipated increases in fire frequency and intensity due to climate change, concomitant with growing population and energy demands, hydroelectric installations and transportation networks in this region face growing risks (Arseneault et al. 2023), emphasizing the need for enhanced fire risk mitigation near these critical infrastructures in remote areas.

3.3. Impacts on wildlife and their habitat: the boreal caribou as a case study

The 2023 Québec wildfires are expected to impact wildlife differently depending on species: burn-associated species (Boucher et al. 2012, 2016) or species adapted to early seral or open forests may benefit (Hutto and Patterson 2016; Knaggs et al. 2020), while those reliant on old-growth forests could suffer due to habitat loss exacerbated by industrial forestry and climate change (Drapeau et al. 2016; Bergeron et al. 2017; Rudolph et al. 2017; Tremblay et al. 2018). In Québec, boreal populations of woodland caribou (*Rangifer tarandus*; hereafter caribou) are respectively listed as threatened and vulnerable under the federal and provincial species at risk acts. Cutovers originating from industrial timber harvesting and associated road networks are regarded as the main causes of caribou population declines in the province (MFFP 2021). Environment Canada (2011) showed that caribou demography was best explained by a combination of wildfire and anthropogenic disturbances, with populations living in heavily disturbed ranges having a lower recruitment rate, and to a lesser extent, a lower adult survival (Johnson et al. 2020). Considering the uniqueness of the 2023 fires, we sought to determine how much of the caribou's distribution range had burned, and we explored the contribution of recent fires on the area covered by total disturbances for several caribou ranges.

Across the 11 caribou ranges studied (Fig. 6), the 2023 wildfires burned 1 490 100 ha, representing on average $2.6\% \pm 3.7\%$ (SD) of range areas (Fig. 7a). The three ranges occurring in northwestern Québec had the highest proportion of area burned, with the Nottaway range showing the highest proportion (11.3%). Before 2023, caribou ranges had on average 43.9% ($\pm 27.1\%$) of their area covered by total disturbances, defined as the sum of natural disturbances (0–40 years old burned areas) and non-overlapping anthropogenic disturbances (0–50 year old clearcuts and roads buffered by 500 m). Most human disturbances occurred south of the northern forest allocation limit, i.e., within the commercial forests. After the 2023 wildfires, caribou ranges were disturbed at 48.0% ($\pm 26.8\%$) on average, representing a 14.0% (± 22.7) increase (i.e., rate of change) compared to pre-2023 conditions (Fig. 7b). Some fires occurred in young regenerating forests, which were already considered as disturbed caribou habitat. These fires had the consequence of burning the regeneration, bringing the areas back to a younger disturbed state, but they did not add to the percent total disturbances in our analyses.

Wildfires in 2023 increased total disturbances in 8 of 11 caribou ranges in Québec. These fires brought the Detour and Nottaway ranges above the 35% management threshold set by


Environment Canada (2012), i.e., the threshold above which a population has less than a 60% probability of being self-sustaining in the long term (Fig. 7b). Before the 2023 fires, 5 of the 11 ranges were already above this threshold. Studies by Environment Canada (2011) and Johnson et al. (2020) have indicated that wildfire impacts on caribou were less severe than those of human disturbances. Caribou are adapted to the dynamic boreal ecosystem, regularly affected by forest fires of varying frequencies and severities (Lafontaine et al. 2019). Therefore, the 2023 fires' additional impacts on caribou were likely modest province-wide, although some populations were more affected than others. Should salvage logging operations be initiated within the distribution range of caribou, the consequences could be detrimental for the species, notably if new permanent roads are introduced into the landscape (Lindenmayer and Noss 2006). Additionally, the cutblocks that would be created following salvage logging would exert a more severe negative impact on caribou populations compared to burned areas (Johnson et al. 2020). In this context, on 8 November 2023, the Québec's Ministry of Natural Resources and Forests prohibited salvage logging operations within a 16 700 ha fire located within the Pimpuacan caribou range overlapping a projected protected area under the stewardship of the Pessamit Innu First Nation.

A sound management and conservation strategy for caribou is urgently needed in Québec. This strategy should implicitly consider the impacts of fire seasons such as that of 2023, expected to occur more frequently in the future (Boulanger et al. 2014). It should also tackle the main cause of caribou declines, i.e., landscape disturbances originating from industrial timber harvesting (St-Laurent et al. 2022; Morineau et al. 2023). Conservation approaches focusing on habitat protection (Leblond et al. 2022) and restoration (Lacerte et al. 2021, 2022) should benefit caribou, as well as other species of the boreal forest (Bichet et al. 2016), although complementary conservation actions would be needed to cover a broader extent of species (Micheletti et al. 2023).

3.4. Impacts on First Nation communities, territories, and people

First Nation communities have historically faced disproportionate consequences of wildfire; many are located within the wildland–urban interface, are exposed to greater fire risks, and are overrepresented in wildfire-related evacuation events (Erni et al. 2021). This trend actualized in 2023, with Eeyou (Cree), Anishnaabe (Algonquin), Atikamekw, and Innu communities experiencing the consequences of wildfires. According to primary data collected by W8banaki emergency center, who have supported First Nations during the 2023 fire season, more than 10 000 people within 13 communities were evacuated. Threats were both direct and indirect and included risks for people and infrastructures, bad air quality over extended periods of time, and closures of roads that were often the only way of accessing the communities. The communities' vulnerability exposed in 2023 is consistent with previous assessments showing that the transition of the Eeyouch from a traditional lifestyle based on trapping, hunt-

Fig. 6. Location of the 11 boreal caribou ranges (black polygons) (A: Detour; B: Nottaway; C: Assinica; D: Témiscamie; E: Caniapiscau; F: Outardes; G: Manicouagan; H: Basse-Côte-Nord; I: Pipmuacan; J: Val d'Or; K: Charlevoix). The map also shows, in orange: the area covered by total disturbances (natural and anthropogenic) as identified in 2020 using the most recent 1:20 000 ecoforest maps published by the Québec government; in red: the 2023 wildfires. Natural disturbances include fires (0–40 years old). Anthropogenic disturbances include roads and clearcuts (0–50 years old), buffered by 500 m. The northern forest allocation limit above which commercial timber harvesting is prohibited is shown using a green line. This figure was created using ArcGIS version 10.8 and assembled from the following data sources: contours of the caribou ranges (Données Québec: <http://www.donneesquebec.ca/recherche/dataset/aires-de-repartition-des-populations-de-caribous-forestier>), ecoforest maps with disturbances (Données Québec: <https://www.donneesquebec.ca/recherche/dataset/carte-ecoforestiere-avec-perturbations>), and the northern forest allocation limit (Données Québec: <https://www.donneesquebec.ca/recherche/dataset/limite-territoriale-des-forets-attribuables>). Base map from GeoGratis (<https://geogratis.gc.ca/>).



ing, fishing, and gathering to a more sedentary way of life has made their new infrastructures more vulnerable to fire (Arseneault et al. 2023).


The wildfires of 2023 have also dramatically transformed First Nation territories. The Eeyou and Anishnaabe family hunting grounds, located in the western boreal zone, were especially affected. Of the 384 Eeyou, Abitibiwiinni, and Lac Simon hunting grounds, 119 (31%) had more than 5% of their land surface burned in 2023. Thirty-one hunting grounds had more than 50% and nine had more than 80%. Hunting grounds are land units passed down from one generation to the next that are key places for cultural and subsistence practices, including trapping, hunting, fishing, speaking language, learning bush skills, and celebrating important life events. The losses experienced by numerous families were thus tremendous, both economically and culturally.

The effects of the 2023 wildfires on First Nation territories accumulate with previous (and future) disturbances from industrial activities, mostly through forestry, mining, and

hydroelectric development (Bélisle and Asselin 2021). For instance, 43 hunting grounds were affected both by 2023 wildfires and by timber harvesting (1990–2020) and are likely to be subjected to salvage logging. As salvage logging operations stem from special plans, they often involve derogations that allow operations to deviate from existing regulations regarding forest roads or the protection of regeneration, for instance, and do not need to undergo the regular public consultation process. Consequently, consultations with First Nations and other regional and local stakeholders are rushed, with only a few weeks to assess and anticipate the impacts of thousands of hectares of salvage logging.

First Nation people have been coping and adapting to wildfire for centuries, developing finely tuned knowledge of burned forests and their resources (Miller et al. 2010). However, for highly disturbed hunting grounds, adding wildfire to previous changes initiated by industrial activities may surpass people's adaptation capacity and affect their quality of life and general wellbeing (Parlee et al. 2012; Fuentes et al.

Fig. 7. (a) Area (in km², gray bars) and percentage (numbers above bars) of boreal caribou ranges that were burned by the 2023 wildfires. (b) Histogram synthesizing the total disturbance levels in boreal caribou ranges before (light gray) and after (dark gray) the 2023 wildfires. The dashed line represents the 35% disturbance management threshold used by Environment and Climate Change Canada to identify populations more likely than not to be self-sustaining in the long term. Percentages above the histogram bars represent the rate of change in total disturbances caused by the 2023 wildfires (i.e., the newly impacted areas), compared to the pre-2023 disturbance levels. *These ranges extend to Ontario; only the Québec portion was analyzed.



2020). Even for hunting grounds with low previous disturbance levels, salvage logging could cause rapid and cascading changes, including increased traffic, industrial development, and land use conflicts (Walker et al. 2011; Bélisle and Asselin 2021). Moreover, the rigid land boundaries between family hunting grounds that were established within the James Bay and Northern Québec Agreement has reduced families' adaptability to large scale disturbances by limiting their access to the entire land (Sénécal and Égré 1999). The interplay between industrial activities and climate change is significantly reshaping the environmental and cultural landscapes in which the First Nation people live, and the extreme fires of 2023 are set to become a defining part of this transformation.

4. Where do we go from here?

The Québec forest industry has faced several crises over the years, including the severe spruce budworm outbreak of the 1970s and 1980s, the softwood lumber crisis (2005–2010), and the *Commission d'étude sur la gestion de la forêt publique Québécoise* (2004). Each time, the industry had to adjust its practices and adapt to the evolving situation. Likewise, Québec has faced important climate-related catastrophes in the last decades (e.g., the 1998 ice storm, the major floods of 1996, 2011, 2017, 2019, and 2020; see the list of all natural disasters in the Canadian Disaster Database; *Public Safety Canada* 2023) that mandated profound changes to the way society operates (for example, see the "*Plan de Protection du Territoire Face aux Inondations*" developed by the Québec government in April 2020 in response to the 2019 flood event, (*Gouvernement du Québec* 2020)). The 2023 wildfires are among these major crises that will pose a substantial challenge for the upcoming years, highlighting the urgency to adapt and reduce the risks associated with future fires. We propose some actions to take below.

Action 1: Considering fire a priori in the calculation of the AAC

When considering overall impacts on timber supply and biodiversity, current forest management in Québec is maladapted to face climate change. Harvesting rates may be too high to sustain a steady timber supply, especially under the new fire regime. One of the most effective and immediate large-scale adaptation strategies would be to create precautionary wood reserves. As a corollary, this implies factoring-in wildfire impacts upfront when calculating the AAC. In Québec, fire effects are typically factored into the AAC a posteriori (with very few exceptions). This means that after a fire occurs, the AAC of the affected forest management unit is recalculated to account for the long-term changes in the availability of harvestable stands caused by fires. In accordance with this strategy, Québec's Forester in Chief has recommended in November 2023, a 12.7%, 2.1%, and 0.2% decrease in AAC within the three regions most affected by the 2023 wildfires (Nord-du-Québec, Abitibi-Témiscamingue, and Mauricie) (*Forestier en Chef* 2023). However, many studies have shown that such a strategy could result in important

variations in AAC over time and would impede sustainability in the long term (Raulier et al. 2014; Leduc et al. 2015; Daniel et al. 2017; *Forestier en Chef* 2021). This prevents the industry from benefiting from predictability in the AAC to plan future activities. Depending on the current and future fire regimes, an a priori consideration of wildfires could lead to the establishment of precautionary wood reserves, i.e., upfront decreases in AAC of e.g., ~5%–20% depending on the probability of having unsustainable timber supply given the fire activity. Although this could initially be seen by some as a negative measure, it would in fact increase the probability of maintaining a constant long-term timber supply predictability, which would be beneficial for the industry in general (Savage et al. 2010). Considering precautionary reserves upfront in the calculation of the AAC would help prevent shortfalls and a posteriori reduction in timber supply for regional burning rates as low as 0.30%–0.45% (Savage et al. 2010; *Ministère des Ressources Naturelles du Québec* 2013). A preliminary analysis conducted for the current study (Suppl. Mat. S3) shows that, if a 20% precautionary reserve had been established 20 years ago in northwestern Québec, the 2023 wildfires would have not led to any drastic postfire decreases in timber supply. Conversely, such a precautionary reserve would have led to more timber harvested in the medium to long term (>2030–2040). Burning rates that would require the establishment of precautionary reserves are already affecting vast areas of the commercial boreal forest in Québec. By mid-century, even under moderate climate change, these rates are expected to impact nearly all the managed boreal forests in Québec (Boulanger et al. 2014; Pau et al. 2023), adding to the urgency of establishing these reserves. Maintaining precautionary wood reserves would also benefit several additional ecosystem services, notably by maintaining old-growth forests that are high-quality habitat for many wildlife species including caribou (Bichet et al. 2016; Leblond et al. 2022; St-Laurent et al. 2022; Labadie et al. 2023).

Action 2: Making forest landscapes more resilient to fire

Increasing forest resilience to wildfires could help reduce postfire regeneration failures. Resilience here refers to the capacity of the forest ecosystem to recover its ecological functions after the disturbance, in this case, wildfires. This could be achieved notably by favoring species with an early sexual maturity such as jack pine (Rudolph and Laidly 1990; Cyr et al. 2022). Variable retention or partial harvesting leaving mature trees after logging could also be envisioned in fire-prone black spruce–jack pine-dominated landscapes; spared trees in a dispersed or aggregated pattern could serve as seeding trees if a burn was to occur a few decades after logging (Perrault-Hébert et al. 2017; Cyr et al. 2022). Perrault-Hébert et al. (2017) showed that leaving between 10% and 15% of mature seed trees could be sufficient to restore a low to moderate level of regeneration and avoid the high social costs of postfire plantations (5–8 k\$.ha⁻¹, when including field preparation and plantation per se, based on 2023 estimations). Seed tree retention could also mitigate the impacts of severe mature biomass

removal on forest biodiversity (Thorn et al. 2020). The precautionary wood reserves discussed above would also mitigate the increase in regeneration failures by leaving more mature stands at the landscape scale. As fires are promoting hardwood species, there is also a need to reevaluate postfire forest management strategies that favor coniferous species over time. Increasing functional redundancy, i.e., “communities with both a mixture of traits that enable species to adapt to known stressors” (Messier et al. 2019), through specific forest management and silvicultural practices, could also help increase forest resilience after disturbance and foster the provision of ecosystem services.

Action 3: Making forest landscapes more resistant to fire

An additional strategy would aim at making forest landscapes more resistant to climate-induced increases in wildfire activity and to reduce their consequences. As opposed to resilience, resistance here refers to the ability of the forest ecosystem to withstand wildfires, notably by being less flammable. Higher resistance could stem from changing, either actively or passively, the flammability of the vegetation, notably through increasing the hardwood component of forest landscapes (Terrier et al. 2013). When fully leafed, hardwood species such as aspen, white birch, and red maple are known to be less flammable than conifer species such as balsam fir, black spruce, and jack pine (Forestry Canada Fire Danger Group 1992; Hély et al. 2001, 2010; Bernier et al. 2016). Simulations have shown that actively planting or favoring the natural regeneration of hardwood species after fire or harvest could strongly alleviate the climate-induced increases in fire activity and mitigate concomitant losses in timber supply by 50% (Forestier en Chef 2021). Preliminary analyses (Suppl. Mat. S4) revealed that hardwood stands, more than any other types of forest cover, were under-represented in the forested areas burned in 2023, underscoring the potential protective qualities of hardwood stands even during severe wildfire seasons.

However, there are several drawbacks to this strategy. The protection ability of hardwood trees is limited before leafout, a period during which a significant number of fires can occur within the boreal forest (Parisien et al. 2023). Furthermore, not all boreal sites can support hardwood species due to specific soil characteristics (Marchais et al. 2022). In the western boreal bioclimatic domain of Québec, hardwood and mixed stands occupy just under 10% of the forest area, primarily on hillsides (Blouin and Berger 2004). The protection ability of hardwood should thus be assessed in the light of the ecological classification of Québec’s forest ecosystems (Saucier et al. 2010). In addition, the operational capacity of the forest sector to convert forest landscapes is limited in space and time. In this context, it might be advantageous to consider how the 2023 fires will themselves alter forest landscapes by increasing the pioneer hardwood components in the forests (Boucher et al. 2014, 2017b).

In addition to these drawbacks, a rapid anthropogenic conversion of boreal landscapes would have tremendously deleterious impacts on a myriad of species associated with conifer

forest covers (Tremblay et al. 2018; Labadie et al. 2023), species that are also typically vulnerable to logging (Imbeau et al. 2001; Venier et al. 2014; Leblond et al. 2022), and climate change (Bouderbala et al. 2023). Rapid conversion of forest landscapes could also significantly alter the livelihoods, cultures, and identities of First Nations who are closely tied to the land (Belisle et al. 2022). In this context, such a strategy could be limited to local conversion of forest landscapes by aiming to decrease wildfire risks to communities or critical infrastructures.

Alternatively, the valuation of wetlands as fire breaks and biological refuges through conservation has been exposed as a contributor to forest resistance and resilience to wildfire in western United States (Fairfax and Whittle 2020). Wetlands provide several ecosystem services (Cimon-Morin et al. 2016) and are of primary importance to culture and subsistence for First Nations, particularly for hunting (Grant 2024), as well as being a passive and inexpensive method of increasing the fire resistance of forests. This unexplored solution could be an opportunity for further collaborative research on wildfires on First Nation lands. However, it is important to note that vegetated wetlands may also carry fire during drought conditions (Canadian Forest Service Fire Danger Group 2021), which are expected to become more prevalent with climate change.

Action 4: Adapting the forest management system and the industrial structure

The 2023 fire season raises questions about the forest sector’s ability to adapt to extreme fire events (Boulanger et al. 2023). Forestry practices, which are based on ecological classifications such as potential vegetations, will certainly have to be revised notably by taking into account the impact of climate change on postfire successional pathways at each ecological classification level (Grondin et al. 2022). On the operational side, the ability of the forest sector to intervene on the landscape is limited. For instance, it might be difficult to restore all the current and future postfire regeneration failures, due to limitations in budget, labor, or nursery capacity. Coping with this situation will need much more financial and operational investments, or we will have to acknowledge that forest productivity could strongly decline. Moreover, the capacity of the forest sector to salvage wood, despite its special plans, will remain logistically limited, and a significant proportion of the burned stands will be left without active management. This underscores the importance of proactively preparing the forest to withstand more frequent fires. The large area burned in 2023 might also represent the opportunity to test alternative management strategies to increase the resistance and resilience of forest landscapes to wildfires and climate change impacts.

Adapting the industrial structure will be paramount to make the whole forest sector more resilient. For example, a significant increase in hardwood content (i.e., aspen and white birch) within the timber supply following increased fire activity might prompt a significant paradigm shift in a forest sector that traditionally relies mostly on conifer species (Brecka et al. 2020). Increased fire activity could lead

to novel uses of salvaged wood, such as pulpwood or bioenergy. Adapting the industrial structure might be more efficient than reactively adapting the forest to rising fire activity, although this will imply developing skills and capacity for action within communities. Innovation in this regard will be crucial, and incentives to promote it will have to be prioritized. In any case, it is likely that the industrial sector will need to focus on anticipating these changes, rather than merely reacting to them, to enhance its own vitality as well as that of the forest communities.

Action 5: Increasing suppression capacity and mitigating risks to communities and infrastructures

Even after considering all of the actions proposed here, the fire management agency's (SOPFEU) operational capacity of 30–40 active fires per day will likely have to be increased if we are to reduce the number of fires that are freely burning each year. On 14 November 2023, the government of Québec announced a 16 mol/L\$ investment in SOPFEU, for 2023–2024, to support increased suppression and prevention capacity (SOPFEU 2023b). This is a step in the right direction, but it may not suffice. Aerial suppression capacity seems to have culminated at just over 3000 h per fire season, as proven by the fire seasons of 2005 and 2023. This could be due to the worldwide aging fleet (Radio-Canada 2023) of the airtankers that are most efficient in our boreal conditions (mainly CL-215 and CL-415; McFayden et al. 2023) and the scarcity of qualified pilots (Noovo Info 2023). New aircrafts would be welcome, and governments may need to commit to purchasing more aircrafts to ensure that they are ready for service by 2030 (Le Soleil 2023). In parallel, new pilots and mechanics will also need to be recruited and trained.

This fire season, lots of resources (staff and machinery) that could have otherwise been tasked to suppression operations were requested to support the fast and reactive responses to safeguard communities and infrastructures from the wildfires, highlighting the essential need for more proactive risk management near sensitive areas. This includes considering risk assessments to identify areas likely to burn and implement mitigation measures around corresponding communities and infrastructures before an emergency occurs. It would also enhance the level of awareness and preparedness among communities in addressing wildfire-related emergencies. For example, antecedent mapping of burn probabilities in the La Grande Rivière hydroelectric complex in the James Bay area could identify the areas that preferentially burned in 2023 (Arseneault et al. 2023), demonstrating the accuracy and usefulness of such predictive assessments in wildfire management (Fig. 8; Parisien et al. 2019). Maintaining an organized database that records the impacts of wildfires on infrastructures is also vital. Such a database would provide valuable insights into the extent and severity of damage and the effectiveness of existing mitigation strategies. This could inform future risk analyses by providing data to develop susceptibility functions for resources and assets threatened by fire, adding to those developed for residential structures (Abo El Ezz et al. 2022; Nicoletta et al. 2023). Furthermore, it is impor-

tant to increase the monitoring capacity, encompassing aspects like evacuation procedures and the resistance of infrastructures. Enhanced monitoring (e.g., through daily remote sensing) would facilitate timely responses during wildfire emergencies, potentially saving lives and reducing property damage. The adoption of FireSmart practices (CIFFC 2023) is a key part of this strategy. In this regard, communities that have established fire breaks in a hurry this year, are now faced with crucial decisions regarding the future of these protective measures. A potential solution could involve intensifying land management near communities, for example, by planting less flammable, fast-growing tree species such as hybrid poplar and hybrid larch, or shrub species like willow and alder, which could also serve purposes like biofuel production (Mansuy et al. 2018). Such measures not only safeguard against the immediate threats of wildfires but also contribute to the long-term resilience and sustainability of these regions. Cognizant of the need for more organized wildland fire risk management, SOPFEU, right after the fire season 2022, created a risk mitigation branch, while the *Ministère de la Sécurité publique du Québec* recently announced 31 mol/L\$ to implement mitigation measures such as fuel treatment and FireSmart measures in communities at high risk of fires.


First Nation communities had to mobilize very quickly to respond to the forest fires of 2023. They updated their emergency response protocols and developed their knowledge of fire risk assessment and fire behavior. They gathered information daily to make decisions with far-reaching consequences for the health and safety of their members. The people who were at the heart of the crisis in the communities were key players in the development of risk assessment and management tools for future fires. In this way, the 2023 fire season may have generated an opportunity for collaboration between scientific and First Nation institutions, both of which possess complementary knowledge and skills.

Action 6: A call for a unified risk management approach

The 2023 forest fires in Québec emphasize the need for a reassessment of risk management strategies considering ongoing climate change and growing systemic risk (Global Assessment Report by UNDRR 2022). It is important to consider risk as a composite notion, made up on the one hand by natural hazards, which are related to the natural dynamics of ecosystems and are directly impacted by global warming, and made up on the other hand by all the factors related to human activities and the development of societies. (IPCC 2018, 2023; UNDRR 2019, 2022). Indeed, focusing risk reduction strategies solely on hazard mitigation, through fuel management for example, without integrating societal parameters as key drivers of fire exposure and vulnerability, could not be sufficient to balance the expected impacts of climate change on fire occurrence, duration, and severity/extent (Barnes et al. 2023; Boulanger et al. 2023), and associated risks (Berry et al. 2010; IPCC 2012; Disse et al. 2020).

Canada has seen significant demographic growth in the last years, particularly in Québec, with a record annual

Fig. 8. Comparison of areas burned in 2023 with fire probabilities mapped in 2022 across the strategic La Grande hydroelectric complex in the James Bay area. Map adapted from [Arseneault et al. \(2023\)](#).



population growth rate of +2.7% in 2022, the highest since 1957, leading to an increase of over 1 million people in 2023 ([Statistics Canada 2023b](#)). This growth trends towards greater community and infrastructure exposure to forest fires, underscoring the urgent need for a multirisk zonation approach to inform municipal and community planning for land use and development. The economic costs associated with natural disasters in Québec, especially forest fires and floods, reached a peak of 738 mol/L\$ in 2023 of exposed and insured assets, a significant increase from averages of around 97 mol/L\$ in 2011–2015 and approximately 222 mol/L\$ in 2016–2020 (CatIQ data, personal communication from B. Marchand). These economic impacts suggest potential growth or closure risks for private insurance properties or industrial assets in the future, raising concerns about socio-economic vulnerability for those within the Intensive Protection Zone and beyond. Additionally, extensive forest fires could cause cascading effects on the water cycle, flood frequency, and environmental health, including impacts on indigenous communities through increased mercury con-

centrations in fish ([Robinne et al. 2018](#); [Garcia and Carignan 2005](#)). Moreover, forest fires could alter flood seasonality and intensity ([Lininger et al. 2021](#); [IPCC 2023](#)). Therefore, this stresses the importance of an integrated risk management approach that holistically considers various climate risks. This will help identify vulnerable communities and ecosystems, and therefore develop integrated adaptation and mitigation strategies to multiple cumulating and interacting risks ([O'Neill et al. 2017](#)).

5. Conclusions

After the 2023 extreme fire season, Québec saw concerted efforts among various stakeholders and First Nations (e.g., *Sommet sur les feux de forêt* ([ORRF 2023](#))) to challenge the untenable status quo and the need for change considering an extreme fire season and ongoing climate changes. A consultation on forest management was initiated by the Québec Minister of Natural Resources and Forests on 15 November 2023, with a view to adapt to this new reality. Current forest

management strategies may no longer be sufficient, requiring a revision of strategies related to the sustainable management of forests, including wood production, protected areas, and wildlife habitats. Such reflections offer an opportunity to update operational methodologies with sustainability principles, particularly for the forest sector, communities, First Nations, and ecosystems.

Our analysis emphasizes that it is crucial to put the impacts and consequences of the 2023 wildfires in the context of increasing fire activity due to climate change. The adoption of the solutions we propose will certainly depend on political decisions and might be difficult to implement without additional resources, either financial or operational. Yet, increasing retention and reducing harvesting rates are actions that could immediately lower the vulnerability of forest landscape to regeneration accidents or timber supply shortages. Favoring deciduous regeneration by reducing treatments that lower their abundance after disturbances is also straightforward. Total societal costs might be way higher under status quo. A continued rise in wildfires under business-as-usual practices could lead to a decline in boreal forest health and its socio-ecological services, impacting the forest industry, carbon sequestration, wildlife and their habitat, and cultural values for indigenous and non-indigenous communities. Some actions could have immediate positive impacts (e.g., Action 5) while others will be felt in the long term. Risk assessment and mitigation, along with adaptation and rapid actions, are key. This includes redefining forestry to make ecosystems and the forest sector more resistant and resilient, identifying vulnerabilities and co-benefits, and implementing regional adaptation measures integrating diverse expertise (Boulanger et al. 2023). Costs of adaptation strategies must be considered, prioritizing approaches with multiple mutual benefits. A precautionary approach is crucial in the face of uncertain climate change (Millar et al. 2007). Systemic risks of forest fires and climate change impacts demand an integrated risk management approach, enhancing preventive tools, and early-warning systems. Experiences like the one we faced during the 2023 fire season are opportunities to remind ourselves that we need to improve fire and forest management strategies and policies. This holistic approach would enhance our ability to predict, prevent, and respond to forest fires, reducing their impacts on economic sectors, ecosystems, and people.

Acknowledgements

We would like to thank Clémence Benoît and the ESCER centre for providing meteorological and FWI data derived from ERA5.

Article information

History dates

Received: 22 December 2023

Accepted: 19 April 2024

Accepted manuscript online: 14 May 2024

Version of record online: 18 December 2024

Copyright

© 2024 Authors Arseneault, Bélisle, Bergeron, Boucher, Danneyrolles, Gachon, Grant, Grondin, Jetté, Labadie, Leduc, and St-Laurent, and The Crown. This work is licensed under a [Creative Commons Attribution 4.0 International License \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Data availability

Data are available upon request to the corresponding author.

Author information

Author ORCIDs

Yan Boulanger <https://orcid.org/0000-0001-6181-8509>

Annie Claude Bélisle <https://orcid.org/0000-0003-1370-2805>

Victor Danneyrolles <https://orcid.org/0000-0002-4839-8164>

Sandy Erni <https://orcid.org/0000-0002-9170-9650>

Eliane Grant <https://orcid.org/0000-0001-7847-0430>

Junior A. Tremblay <https://orcid.org/0000-0003-4930-0939>

Kaysandra Waldron <https://orcid.org/0000-0001-7551-4346>

Author notes

Jean-Pierre Jetté is an independent professional forester.

Author contributions

Conceptualization: YB

Formal analysis: YB, DA, ACB, JB, VD, SE, PG, MPG, ML, AL, JPP, MS

Methodology: YB, DA, ACB, JB, VD, PG, MPG, JJ, ML, AL, MS

Project administration: YB

Supervision: YB

Writing – original draft: YB, DA, ACB, JB, YB, VD, SE, PG, MPG, EG, PG, JJ, GL, ML, AL, MS, JAT, KW

Writing – review & editing: YB, DA, ACB, YB, JB, YB, VD, SE, PG, MPG, EG, PG, JJ, GL, ML, AL, JPP, MS, JAT, KW

Competing interests

The authors declare there are no competing interests.

Supplementary material

Supplementary data are available with the article at <https://doi.org/10.1139/cjfr-2023-0298>.

References

- Abo El Ezz, A., Boucher, J., Cotton-Gagnon, A., and Godbout, A. 2022. Framework for spatial incident-level wildfire risk modelling to residential structures at the wildland urban interface. *Fire Saf. J.* **131**: 103625. doi:[10.1016/j.firesaf.2022.103625](https://doi.org/10.1016/j.firesaf.2022.103625).
- Antokhina, O.Y., Antokhin, P.N., Belan, B.D., Gochakov, A.V., Martynova, Y.V., Pustovalov, K.N., et al. 2023. Effects of Rossby waves breaking and

- atmospheric blocking formation on the extreme forest fire and floods in Eastern Siberia 2019. *Fire*, **6**(3): 122. doi:[10.3390/fire6030122](https://doi.org/10.3390/fire6030122).
- Arseneault, D., Parisien, M.-A., Boulanger, Y., Simard, M., Asselin, H., Shakeri, Z., et al. 2023. Exposition et adaptation aux incendies forestiers dans la taiga canadienne. Scientific Report.
- Barnes, C., Boulanger, Y., Keeping, T., Gachon, P., Gillett, N., Boucher, J., et al. 2023. Climate change more than doubled the likelihood of extreme fire weather conditions in eastern Canada. *Scientific Report*. doi:[10.25561/105981](https://doi.org/10.25561/105981).
- Bélisle, A.C., and Asselin, H. 2021. A collaborative typology of boreal indigenous landscapes. *Can. J. For. Res.* **51**(8): 1–10. doi:[10.1139/cjfr-2020-0369](https://doi.org/10.1139/cjfr-2020-0369).
- Bélisle, A.C., Gauthier, S., and Asselin, H. 2022. Integrating indigenous and scientific perspectives on environmental changes: insights from boreal landscapes. *People Nat.* **4**(6): 1513–1535. doi:[10.1002/PAN3.10399](https://doi.org/10.1002/PAN3.10399).
- Bergeron, Y., Cyr, D., Girardin, M.P., and Carcaillet, C. 2010. Will climate change drive 21st century burn rates in Canadian boreal forest outside of its natural variability: collating global climate model experiments with sedimentary charcoal data. *Int. J. Wildl. Fire*, **19**: 1127–1139. doi:[10.1071/WF09092](https://doi.org/10.1071/WF09092).
- Bergeron, Y., Vijayakumar, D.B.I.P., Ouzennou, H., Raulier, F., Leduc, A., and Gauthier, S. 2017. Projections of future forest age class structure under the influence of fire and harvesting: implications for forest management in the boreal forest of eastern Canada. *Forestry*, **90**(4): 485–495. doi:[10.1093/forestry/cpx022](https://doi.org/10.1093/forestry/cpx022).
- Bernier, P.Y., Gauthier, S., Jean, P.-O., Manka, F., Boulanger, Y., Beaudoin, A., and Guindon, L. 2016. Mapping local effects of forest properties on fire risk across Canada. *Forests*, **7**(8): 157. doi:[10.3390/f7080157](https://doi.org/10.3390/f7080157).
- Berry, H.L., Bowen, K., and Kjellstrom, T. 2010. Climate change and mental health: a causal pathways framework. *Int. J. Public Health*, **55**(2): 123–132. doi:[10.1007/s00038-009-0112-0](https://doi.org/10.1007/s00038-009-0112-0).
- Bichet, O., Dupuch, A., Hébert, C., Le Borgne, H., and Fortin, D. 2016. Maintaining animal assemblages through single-species management: the case of threatened caribou in boreal forest. *Ecol. Appl.* **26**: 612–623. doi:[10.1890/15-0525](https://doi.org/10.1890/15-0525). PMID: 27209799.
- Blouin, J., and Berger, J.P. 2004. Guide de reconnaissance des types écologiques des régions écologiques 6c—Plaine du lac Opémisca, 6d—Coteaux du lac Assinica, 6e—Coteaux de la rivière Nestaoacan, 6f—Coteaux du lac Mistassini et 6 g—Coteaux du lac Manouane, ministère des Ressources naturelles, de la Faune et des Parcs, Forêt Québec, Direction des inventaires forestiers, Division de la classification écologique et productivité des stations, Québec, QC, Canada.
- Boucher, J., Azeria, E.T., Ibarzabal, J., and Hébert, C. 2012. Saproxyllic beetles in disturbed boreal forests: temporal dynamics, habitat associations, and community structure. *Écoscience*, **19**(4): 328–343. doi:[10.2980/19-4-3503](https://doi.org/10.2980/19-4-3503).
- Boucher, J., Hébert, C., Ibarzabal, J., and Bauce, É. 2016. High conservation value forests for burn-associated saproxyllic beetles: an approach for developing sustainable post-fire salvage logging in boreal forest. *Insect Conserv. Divers.* **9**(5): 402–415. doi:[10.1111/icad.12175](https://doi.org/10.1111/icad.12175).
- Boucher, J., Hébert, C., and Bauce, E. 2020. A flexible approach for predicting and mapping postfire wood attacks in black spruce and jack pine forests using the differenced normalized burn ratio (dnBR). *Can. J. For. Res.* **50**: 880–889. doi:[10.1139/cjfr-2019-0334](https://doi.org/10.1139/cjfr-2019-0334).
- Boucher, Y., Grondin, P., and Auger, I. 2014. Land use history (1840–2005) and physiography as determinants of southern boreal forests. *Landsc. Ecol.* **29**: 437–450. doi:[10.1007/s10980-013-9974-x](https://doi.org/10.1007/s10980-013-9974-x).
- Boucher, Y., Perrault-Hébert, M., Fournier, R., Drapeau, P., and Auger, I. 2017a. Cumulative patterns of logging and fire (1940–2009): consequences on the structure of the eastern Canadian boreal forest. *Landsc. Ecol.* **32**(2): 361–375. doi:[10.1007/s10980-016-0448-9](https://doi.org/10.1007/s10980-016-0448-9).
- Boucher, Y., Auger, I., Noël, J., Grondin, P., and Arseneault, D. 2017b. Fire is a stronger driver of forest composition than logging in the boreal forest of eastern Canada. *J. Veg. Sci.* **28**(1): 57–68. doi:[10.1111/jvs.12466](https://doi.org/10.1111/jvs.12466).
- Bouderbala, I., Labadie, G., Béland, J.M., Tremblay, J.A., Boulanger, Y., Hébert, C., et al. 2023. Long-term effect of forest harvesting on boreal species assemblages under climate change. *PLOS Clim.* **2**(3): e0000179. doi:[10.1371/journal.pclm.0000179](https://doi.org/10.1371/journal.pclm.0000179).
- Boulanger, Y., and Pascual Puigdevall, J. 2021. Boreal forests will be more severely affected by projected anthropogenic climate forcing than mixedwood and northern hardwood forests in eastern Canada. *Landsc. Ecol.* **36**: 1725–1740. doi:[10.1007/s10980-021-01241-7](https://doi.org/10.1007/s10980-021-01241-7).
- Boulanger, Y., Gauthier, S., and Burton, P.J. 2014. A refinement of models projecting future Canadian fire regimes using homogeneous fire regime zones. *Can. J. For. Res.* **44**: 365–376. doi:[10.1139/cjfr-2013-0372](https://doi.org/10.1139/cjfr-2013-0372).
- Boulanger, Y., Pascual, J., Bouchard, M., D’Orangeville, L., Périé, C., and Girardin, M.P. 2022a. Multi-model projections of tree species performance in Québec, Canada under future climate change. *Global Change Biol.* **28**: 1884–1902. doi:[10.1111/gcb.16014](https://doi.org/10.1111/gcb.16014).
- Boulanger, Y., Boucher, J., and Pascual Puigdevall, J. 2022b. Climate change impacts on suppression team workload. *Wildland Fire Canada Conference*. October 31–November 4 2022, Edmonton, Canada.
- Boulanger, Y., Puigdevall, J.P., Belisle, A.C., Bergeron, Y., Brice, M.-H., Cyr, D., et al. 2023. A regional integrated assessment of the impacts of climate change and of the potential adaptation avenues for Québec’s forests. *Can. J. For. Res.* **53**(8): 556–578. doi:[10.1139/cjfr-2022-0282](https://doi.org/10.1139/cjfr-2022-0282).
- Brecka, A.F., Boulanger, Y., Searle, E.B., Taylor, A.R., Price, D.T., Zhu, Y., et al. 2020. Sustainability of Canada’s forestry sector may be compromised by impending climate change. *For. Forest Ecol. Manage.* **474**: 118352. doi:[10.1016/j.foreco.2020.118352](https://doi.org/10.1016/j.foreco.2020.118352).
- Canadian Forest Service. 2023. Canadian National Fire Database—Agency Fire Data [online]. Natural Resources Canada, Canadian Forest Service, Northern Forestry Centre, Edmonton, AB. Available from <http://cwfis.cfs.nrcan.gc.ca/ha/nfdb> [accessed 22 October 2023].
- Canadian Forest Service Fire Danger Group. 2021. An overview of the next generation of the Canadian Forest Fire Danger Rating System. Information Report GLC-X-26. Natural Resources Canada, Canadian Forest Service, Great Lakes Forestry Centre. 70p.
- Canadian Interagency Forest Fire Centre [CIFFC]. 2023. Canadian Interagency Forest Fire Centre. Available from ciffc.ca [accessed 22 October 2023].
- Canadian Wildland Fire Information System [CWFFIS]. 2023. The Canadian Wildland Fire Information System. Available from <https://cwfis.cfs.nrcan.gc.ca/> [accessed 22 October 2023].
- Carcaillet, C., Bergeron, Y., Richard, P.J.H., Fréchette, B., Gauthier, S., and Prairie, Y.T. 2001. Change of fire frequency in the eastern Canadian boreal forests during the Holocene: does vegetation composition or climate trigger the fire regime? *J. Ecol.* **89**(6): 930–946. doi:[10.1111/j.1365-2745.2001.00614.x](https://doi.org/10.1111/j.1365-2745.2001.00614.x).
- Cardil, A., Lorente, M., Boucher, D., Boucher, J., and Gauthier, S. 2019. Factors influencing fire suppression success in the province of Québec (Canada). *Can. J. For. Res.* **49**(5): 531–542. doi:[10.1139/cjfr-2018-0272](https://doi.org/10.1139/cjfr-2018-0272).
- CBS News. 2023. New Jersey raises air quality alert to “unhealthy” due to smoke from Canada wildfires. Available from <https://www.cbsnews.com/newyork/news/new-jersey-raises-air-quality-alert-to-unhealthy-due-to-smoke-from-canada-wildfires> [accessed 4 December 2023].
- Chaste, E., Girardin, M.P., Kaplan, J.O., Bergeron, Y., and Hély, C. 2019. Increases in heat-induced tree mortality could drive reductions of biomass resources in Canada’s managed boreal forest. *Landsc. Ecol.* **34**: 403–426.
- Chavardès, R.D., Danneyrolles, V., Portier, J., Girardin, M.P., Gaboriau, D.M., Gauthier, S., et al. 2022. Converging and diverging burn rates in North American boreal forests from the Little Ice Age to the present. *Int. J. Wildl. Fire*, **31**: 1184–1193. doi:[10.1071/WF22090](https://doi.org/10.1071/WF22090).
- Cimon-Morin, J., Darveau, M., and Poulin, M. 2016. Site complementarity between biodiversity and ecosystem services in conservation planning of sparsely-populated regions. *Environ. Conserv.* **43**(1): 56–68. doi:[10.1017/S0376892915000132](https://doi.org/10.1017/S0376892915000132).
- Commission d’étude sur la gestion de la forêt publique québécoise. 2004. Rapport de la Commission d’étude sur la gestion de la forêt publique québécoise. Ministère de la Forêt, de la Faune et des Parcs du Québec. ISBN 2-550-43626-1.
- Cyr, D., Splawinski, T.B., Pascual Puigdevall, J., Valeria, O., Leduc, A., Thiffault, N., et al. 2022. Mitigating post-fire regeneration failure in boreal landscapes with reforestation and variable retention harvesting: at what cost? *Can. J. For. Res.* **52**(4): 568–581. doi:[10.1139/cjfr-2021-0180](https://doi.org/10.1139/cjfr-2021-0180).
- Daniel, C.J., Ter-Mikaelian, M.T., Wotton, B.M., Rayfield, B., and Fortin, M.J. 2017. Incorporating uncertainty into forest management planning: timber harvest, wildfire and climate change in the boreal forest. *Forest Ecol. Manage.* **400**: 542–554. doi:[10.1016/j.foreco.2017.06.039](https://doi.org/10.1016/j.foreco.2017.06.039).

- Danneyyrolles, V., Cyr, D., Girardin, M.P., Gauthier, S., Asselin, H., and Bergeron, Y. 2021. Influences of climate fluctuations on northeastern North America's burned areas largely outweigh those of European settlement since AD 1850. *Environ. Res. Lett.* **16**(11): 114007. doi:10.1088/1748-9326/ac2ce7.
- Danneyyrolles, V., Boucher, Y., Fournier, R., and Valeria, O. 2023. Positive effects of projected climate change on post-disturbance forest regrowth rates in northeastern North American boreal forests. *Environ. Res. Lett.* **18**(2): 024041. doi:10.1088/1748-9326/acb72a.
- Disse, M., Johnson, T.G., Leandro, J., and Hartmann, T. 2020. "Exploring the relation between flood risk management and flood resilience". *Water Secur.* **9**: 100059. doi:10.1016/j.wasec.2020.100059.
- Drapeau, P., Villard, M., Leduc, A., and Hannon, S.J. 2016. Natural disturbance regimes as templates for the response of bird species assemblages to contemporary forest management. *Divers. Distrib.* **22**: 385–399. doi:10.1111/ddi.12407.
- Drobyshev, I., Bergeron, Y., Girardin, M.P., Gauthier, S., Ols, C., and Ojal, J. 2017. Strong gradients in forest sensitivity to climate change revealed by dynamics of forest fire cycles in the post little ice age era: forest fires in post LIA era. *J. Geophys. Res. Biogeosci.* **122**(10): 2605–2616. doi:10.1002/2017JG003826.
- Ellis, T.M., Bowman, D.M.J.S., Jain, P., Flannigan, M.D., and Williamson, G.J. 2022. Global increase in wildfire risk due to climate-driven declines in fuel moisture. *Global Change Biol.* **28**: 1544–1559. doi:10.1111/gcb.16006.
- Environment Canada. 2011. Scientific assessment to inform the identification of critical habitat for woodland caribou (*Rangifer tarandus caribou*), boreal populations, in Canada: 2011 update.
- Environment Canada. 2012. Recovery Strategy for the woodland caribou (*Rangifer tarandus caribou*), boreal population, in Canada. Species at Risk Act Recovery Strategy Series. Environment Canada, Ottawa. xi + 138pp.
- Erni, S., Arseneault, D., Parisien, M.A., and Bégin, Y. 2017. Spatial and temporal dimensions of fire activity in the fire-prone eastern Canadian taiga. *Global Change Biol.* **23**(3): 1152–1166. doi:10.1111/gcb.13461.
- Erni, S., Johnston, L., Boulanger, Y., Manka, F., Bernier, P., Eddy, B., et al. 2021. Exposure of the Canadian wildland–human interface and population to wildland fire, under current and future climate conditions. *Can. J. For. Res.* **51**: 1357–1367. doi:10.1139/cjfr-2020-0422.
- ESCCER. 2023. Centre pour l'étude et la simulation du climat à l'échelle régionale. Trends and climate normals of weather variables and indices. Available from http://feux.escer.uqam.ca/en/we_monitoring.html [accessed 22 October 2023].
- ESRI. 2017. ArcGIS desktop: ArcMap release 10.5.1. Environmental Systems Research Institute, Redlands, CA.
- Fairfax, E., and Whittle, A. 2020. Smokey the beaver: beaver-dammed riparian corridors stay green during wildfire throughout the western United States. *Ecol. Appl.* **30**(8): e02225. doi:10.1002/eap.2225. PMID: 32881199.
- Forestier en chef. 2021. Integration of climate change and development of adaptive capacity in the determination of harvest levels in Québec, Roberval, Québec, Available from: https://forestierenchef.gouv.qc.ca/wp-content/uploads/bfec_cc_rapport_veng_02_02_2023.pdf [accessed 15 December 2023]. p. 60.
- Forestier en chef. 2023. Feux de forêt de l'été 2023—Le Forestier en chef recommande une modification des possibilités forestières pour les régions du Nord-du-Québec, de l'Abitibi-Témiscamingue et de la Mauricie. Available from <https://forestierenchef.gouv.qc.ca/nouvelles-communiques-de-presse/feux-de-foret-de-lete-2023/> [accessed 5 December 2023].
- Forestry Canada Fire Danger Group. 1992. Development and structure of the Canadian Forest Fire Behavior Prediction System. Forestry Canada. Forestry Canada, Headquarters, Fire Danger Group and Science and Sustainable Development Directorate, Ottawa. Information Report ST-X-3. 64p.
- Fuentes, L., Asselin, H., Bélisle, A.C., and Labra, O. 2020. Impacts of environmental changes on well-being in indigenous communities in Eastern Canada. *Int. J. Environ. Res. Public Health*, **17**(2): 637. doi:10.3390/ijerph17020637. PMID: 31963806.
- García, E., and Carignan, R. 2005. Mercury concentrations in fish from forest harvesting and fire-impacted Canadian Boreal lakes compared using stable isotopes of nitrogen. *Environ. Toxicol. Chem.* **24**: 685–693. doi:10.1897/04-065R.1. PMID: 15779770.
- Gauthier, S., Bernier, P., Kuuluvainen, T., Shvidenko, A.Z., and Schepaschenko, D.G. 2015. Boreal forest health and global change. *Science*, **349**(6250): 819–822. doi:10.1126/science.aaa9092. PMID: 26293953.
- Gennaretti, F., Arseneault, D., Nicault, A., Perreault, L., and Bégin, Y. 2014. Volcano-induced regime shifts in millennial tree-ring chronologies from northeastern North America. *Proc. Natl. Acad. Sci. U.S.A.* **111**(28): 10077–10082. doi:10.1073/pnas.1324220111.
- Girardin, M.P., and Wotton, B.M. 2009. Summer moisture and wildfire risks across Canada. *J. Appl. Meteorol. Climatol.* **48**: 517–533. doi:10.1175/2008JAMC1996.1.
- Girardin, M.P., Ali, A.A., Carcaillet, C., Mudelsee, M., Drobyshev, I., Hély, C., and Bergeron, Y. 2009. Heterogeneous response of circumboreal wildfire risk to climate change since the early 1900s. *Global Change Biol.* **15**: 2751–2769. doi:10.1111/j.1365-2486.2009.01869.x.
- Girardin, M.P., Portier, J., Remy, C.C., Ali, A.A., Paillard, J., Blarquez, O., et al. 2019. Coherent signature of warming-induced extreme sub-continental boreal wildfire activity 4800 and 1100 years BP. *Environ. Res. Lett.* **14**(12): 124042. doi:10.1088/1748-9326/ab59c9.
- Gouvernement du Québec. 2020. Plan de Protection du Territoire Face aux Inondations: des solutions durables pour mieux protéger nos milieux de vie; Gouvernement du Québec, Available from https://cdn-contenu.quebec.ca/cdn-contenu/adm/min/affaires-municipales/publications-adm/documents/plan_protection_territoire_inondations/PLA_inondations.pdf [accessed 15 December 2023].
- Gouvernement du Québec. 2023a. Aide financière en raison des feux de forêt. Available from <https://www.quebec.ca/securite-situations-urgence/urgences-sinistres-risques-naturels/obtenir-aide-sinistre/aide-financiere-feux-forets> [accessed 21 December 2023].
- Gouvernement du Québec. 2023b. Données Québec. Baux de villégiature. Available from <https://www.donneesquebec.ca/recherche/dataset/couche-des-droits-fonciers-baux> [accessed 20 December 2023].
- Grant, E. 2023. The consequences of fires for the Cree users in the Eeyou Itstchee territory. [In French] [Conference presentation]. In Twenty Fifth Colloquium of the Chair in Sustainable Forest Management, 28 November 2023, Rouyn-Noranda, Québec, Canada. Available from <https://chaireafd.uqat.ca/evenements/colloque-chaire-2023/colloque-chaire-afd-2023-cahier-participant.pdf> [accessed 18 June 2024].
- Grant, E. 2024. Effets des perturbations anthropiques sur l'utilisation autochtone des milieux humides, particulièrement pour la chasse à l'original. Unpublished master's thesis, Université du Québec en Abitibi-Témiscamingue.
- Grondin, P., Brice, M.H., Boulanger, Y., Morneau, C., Couillard, P.-L., Richard, P.J.H., et al. 2022. Ecological classification in forest ecosystem management: links between current practices and future climate change in a Québec case study. In Chapter 8 in Boreal forests in the face of climate change. Edited by M.M. Girona, H. Morin, S. Gauthier et Y. Bergeron.
- Hély, C., Flannigan, M., Bergeron, Y., and McRae, D. 2001. Role of vegetation and weather on fire behavior in the Canadian mixedwood boreal forest using two fire behavior prediction systems. *Can. J. For. Res.* **31**: 430–441. doi:10.1139/x00-192.
- Hély, C., Fortin, M.-J., Anderson, K.R., and Bergeron, Y. 2010. Landscape composition influences local pattern of fire size in the eastern Canadian boreal forest: role of weather and landscape mosaic on fire size distribution in mixedwood boreal forest using the Prescribed Fire Analysis System. *Int. J. Wildl. Fire*, **19**: 1099–1109. doi:10.1071/WF09112.
- Héon, J., Arseneault, D., and Parisien, M.-A. 2014. Resistance of the boreal forest to high burn rates. *Proc. Natl. Acad. Sci. U.S.A.* **111**(38): 13888–13893. doi:10.1073/pnas.1409316111.
- Hersbach, H., Pell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., et al. 2020. The ERA5 global reanalysis. *Quart. J. R. Meteorol. Soc.* **146**. doi:10.1002/qj.3803.
- Hirsch, K., and Martell, D.A. 1996. Review of initial attack fire crew productivity and effectiveness. *Int. J. Wildl. Fire*, **6**: 199–215. doi:10.1071/WF9960199.
- Hope, E.S., McKenney, D.W., Pedlar, J.H., Stocks, B.J., and Gauthier, S. 2016. Wildfire suppression costs for Canada under a changing climate. *PLoS ONE*, **11**(8): e0157425. doi:10.1371/journal.pone.0157425. PMID: 27513660.

- Hutto, R.L., and Patterson, D.A. 2016. Positive effects of fire on birds may appear only under narrow combinations of fire severity and time-since-fire. *Int. J. Wildl. Fire*, **25**: 1074–1085. doi:10.1071/WF15228.
- Imbeau, L., Monkkonen, M., and Desrochers, A. 2001. Long-term effects of forestry on birds of the eastern Canadian boreal forests: a comparison with Fennoscandia. *Conserv. Biol.* **15**(4): 1151–1162. doi:10.1046/j.1523-1739.2001.0150041151.x.
- IPCC. 2012. Managing the risks of extreme events and disasters to advance climate change adaptation. In *A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change*. Edited by C.B. Field, V. Barros, T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi, M.D. Mastrandrea, K.J. Mach, G.-K. Plattner, S.K. Allen, M. Tignor and P.M. Midgley. Cambridge University Press, Cambridge, UK, and New York, NY, USA. 582p.
- IPCC. 2018. Summary for policymakers. In *Global Warming of 1.5° C. An IPCC Special Report on the Impacts of Global Warming of 1.5° C above Pre-industrial Levels and Related Global Greenhouse Gas Emission Pathways, in the Context of Strengthening the Global Response to the Threat of Climate Change, Sustainable Development, and Efforts to Eradicate Poverty*. Edited by V. Masson-Delmotte, P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J.B.R. Matthews, Y. Chen, X. Zhou, M.I. Gomis, E. Lonnoy, T. Maycock, M. Tignor and T. Waterfield. Cambridge University Press, Cambridge, UK and New York, NY, USA. 616p. doi:10.1017/9781009157940.
- IPCC. 2023. Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change Edited by H. Lee and J. Romero. IPCC, Geneva, Switzerland. 184p. doi:10.59327/IPCC/AR6-9789291691647.
- Jain, P., Wang, X., and Flannigan, M.D. 2017. Trend analysis of fire season length and extreme fire weather in North America between 1979 and 2015. *Int. J. Wildl. Fire*, **26**: 1009–1020. doi:10.1071/WF17008.
- Jain, P., Castellanos-Acuna, D., Coogan, S.C.P., Abatzoglou, J.T., and Flannigan, M.D. 2022. Observed increases in extreme fire weather driven by atmospheric humidity and temperature. *Nat. Clim. Change*, **12**: 63–70. doi:10.1038/s41558-021-01224-1.
- Johnson, C.A., Sutherland, G.D., Neave, E., Leblond, M., Kirby, P., Superbie, C., and McLoughlin, P.D. 2020. Science to inform policy: linking population dynamics to habitat for a threatened species in Canada. *J. Appl. Ecol.* **57**: 1314–1327. doi:10.1111/1365-2664.13637.
- Knaggs, M., Haché, S., Nielsen, S.E., Pankratz, R.F., and Bayne, E. 2020. Avian response to wildfire severity in a northern boreal region. *Forests*, **11**: 1330. doi:10.3390/f11121330.
- La Presse. 2023. Épuisée, la mairesse de Chapais démissionne. LaPresse. Available from <https://www.lapresse.ca/actualites/regional/2023-11-08/nord-du-Québec/epuisee-la-mairesse-de-chapais-demissionne.php> [accessed 18 December 2023].
- Labadie, G., Bouderbala, I., Boulanger, Y., Béland, J.M., Hébert, C., Allard, A., et al. 2023. The umbrella value of caribou management strategies for biodiversity conservation in boreal forests under global change. *Sci. Total Environ.* **907**: 168087. doi:10.1016/j.scitotenv.2023.168087. PMID: 37879475.
- Lacerte, R., Leblond, M., and St-Laurent, M.-H. 2021. Determinants of vegetation regeneration on forest roads following restoration treatments: implications for boreal caribou conservation. *Restor. Ecol.* **29**(7): e13414. doi:10.1111/rec.13414.
- Lacerte, R., Leblond, M., and St-Laurent, M.-H. 2022. End of the road: short-term responses of a large mammal community to forest road decommissioning. *J. Nat. Conserv.* **69**: 126256. doi:10.1016/j.jnc.2022.126256.
- Lafontaine, A., Drapeau, P., Fortin, D., Gauthier, S., Boulanger, Y., and St-Laurent, M.-H. 2019. Exposure to historical burn rates shapes the response of boreal caribou to timber harvesting. *Ecosphere*, **10**: e02739. doi:10.1002/ecs2.2739.
- Le Devoir. 2023a. De vastes pans de la forêt boréale n'arriveront pas à se régénérer. Available from <https://www.ledevoir.com/environnement/796885/de-vastes-pans-de-la-foret-boreale-Quebecoise-n-arriveront-pas-a-se-regenerer> [accessed 22 October 2023].
- Le Devoir. 2023b. Les feux de forêt causent de nouvelles pannes d'électricité dans la région de Montréal. Available <https://www.ledevoir.com/societe/794515/les-feux-de-forets-causent-de-nouvelles-pannes-d-electricite-dans-la-region-de-montreal>. [accessed 18 December 2023].
- Le Soleil. 2023. Avions-citernes: des experts déplorent une perte d'expertise au pays. Available from <https://www.lesoleil.com/actualites/2023/06/06/avions-citernes-des-experts-deplorent-une-perde-d-expertise-au-pays-INFTVNYJFZGIDPMWRK12AYR2WU/> [accessed 21 December 2023].
- Leblond, M., Rudolph, T., Boisjoly, D., Dussault, C., and St-Laurent, M.-H. 2022. Science-informed policy decisions lead to the creation of a protected area for a wide-ranging species at risk. *Conserv. Sci. Pract.* **4**: e12833. doi:10.1111/CSP2.12833.
- Leduc, A., Bernier, P.Y., Mansuy, N., Raulier, F., Gauthier, S., and Bergeron, Y. 2015. Using salvage logging and tolerance to risk to reduce the impact of forest fires on timber supply calculations. *Can. J. For. Res.* **45**: 480–486. doi:10.1139/cjfr-2014-0434.
- Leibniz Institute for Tropospheric Research. 2023. Smoke from Canadian wildfires has been hanging over Germany for weeks. Available from <https://www.tropos.de/en/current-issues/press-releases/details/rauch-von-kanadischen-waldbraenden-schwebt-seit-wochen-ueber-deutschland> [accessed 4 December 2023].
- Leverkus, A.B., Benayas, J.M.R., Castro, J., Boucher, D., Brewer, S., Collins, B.M., et al. 2018. Salvage logging effects on regulating and supporting ecosystem services—a systematic map. *Can. J. For. Res.* **48**(9): 983–1000. doi:10.1139/cjfr-2018-0114.
- Leverkus, A.B., Gustafsson, L., Lindenmayer, D.B., Castro, J., Rey Benayas, J.M., Ranius, T., and Thorn, S. 2020. Salvage logging effects on regulating ecosystem services and fuel loads. *Front. Ecol. Environ.* **18**(7): 391–400. doi:10.1002/fee.2219.
- Lindenmayer, D.B., and Noss, R.F. 2006. Salvage logging, ecosystem processes, and biodiversity conservation. *Conserv. Biol.* **20**(4): 949–958. doi:10.1111/j.1523-1739.2006.00497.x. PMID: 16922212.
- Lininger, K.B., Scamardo, J.E., and Guiney, M.R. 2021. Floodplain large wood and organic matter jam formation after a large flood: investigating the influence of floodplain forest stand characteristics and river corridor morphology. *J. Geophys. Res. Earth Surf.* **126**: e2020JF006011. doi:10.1029/2020JF006011.
- Mansuy, N., Barrette, J., Laganière, J., Mabee, W., Paré, D., Gautam, S., et al. 2018. Salvage harvesting for bioenergy in Canada: from sustainable and integrated supply chain to climate change mitigation. *WIREs Energy Environ.* **7**: 298. doi:10.1002/wene.298.
- Marchais, M., Arseneault, D., and Bergeron, Y. 2022. The rapid expansion of *Populus tremuloides* due to anthropogenic disturbances in eastern Canada. *Can. J. For. Res.* **52**(7): 991–1001. doi:10.1139/cjfr-2022-0082.
- McFayden, C.B., Wotton, B.M., Robinson, J.W., Johnston, J.M., Cantin, A., Jurko, N.M., et al. 2023. Reference guide to the drop effectiveness of skimmer and rotary wing airtankers. Information Report GLC-X-35. Natural Resources Canada, Canadian Forest Service. Great Lakes Forestry Centre, Sault Ste. Marie, Ontario. 182p.
- Messier, C., Bauhus, J., Doyon, F., Maure, F., Sous-Silva, R., Nolet, P., et al. 2019. The functional complex network approach to foster forest resilience to global changes. *Forest Ecosyst.* **6**: 21. doi:10.1186/s40663-019-0166-2.
- Micheletti, T., Haché, S., Stralberg, D., Stewart, F.E.C., Chubaty, A.M., Barros, C., et al. 2023. Will this umbrella leak? A caribou umbrella index for boreal landbird conservation. *Conserv. Sci. Pract.* **5**: e12908. doi:10.1111/csp2.12908.
- Microsoft Canadian Buildings Footprint [MCBF]. 2020. GitHub, Canadian buildings footprint, Microsoft. Available from <https://github.com/microsoft/CanadianBuildingFootprints> [accessed 20 December 2023].
- Millar, C., Stephenson, N.L., and Stephens, S.L. 2007. Climate change and forests of the future: managing in the face of uncertainty. *Ecol. Appl.* **17**: 2145–2151. doi:10.1890/06-1715.1. PMID: 18213958.
- Miller, A.M., Davidson-Hunt, I.J., and Peters, P. 2010. Talking about fire: Pikangikum First Nation elders guiding fire management. *Can. J. For. Res.* **40**(12): 2290–2301. doi:10.1139/X10-177.
- Ministère de la Forêt, de la Faune et des Parcs [MFFP]. 2021. Revue de littérature sur les facteurs impliqués dans le déclin des populations de caribous forestiers au Québec et de caribous montagnards de la Gaspésie. Gouvernement du Québec, Québec. pp. xix + 260.
- Ministère des Ressources naturelles du Québec. 2013. Rapport du Comité scientifique chargé d'examiner la limite nordique des forêts attribuables. Secteur des forêts. 148p. + 6 annexes.

- Ministère des Ressources naturelles du Québec. 2019. La Base de données géographiques et administratives à l'échelle de 1/1 000 000. Available from https://mern.gouv.qc.ca/publications/territoire/portrait/1_M/energ_l.zip [accessed 18 June 2024].
- Morineau, C., Boulanger, Y., Gachon, P., Plante, S., and St-Laurent, M.-H. 2023. Climate change alone cannot explain boreal caribou range recession in Québec since 1850. *Global Change Biol.* **29**: 6661–6678. doi:10.1111/gcb.16949.
- Mudelsee, M. 2002. XTREND: a computer program for estimating trends in the occurrence rate of extreme weather and climate events *In* Scientific Reports. Vol. 26. Edited by A. Raabe and K. Arnold. Institute of Meteorology, University of Leipzig. pp. 149–195.
- Mudelsee, M., Börngen, M., Tetzlaff, G., and Grünewald, U. 2004. Extreme floods in central Europe over the past 500 years: role of cyclone pathway “Zugstrasse Vb”. *J. Geophys. Res. Atmos.* **109**(D23): D23101. doi:10.1029/2004JD005034.
- Nappi, A., and Drapeau, P. 2011. Pre-fire forest conditions and fire severity as determinants of the quality of burned forests for deadwood-dependent species: the case of the black-backed woodpecker. *Can. J. For. Res.* **41**: 994–1003. doi:10.1139/x11-028.
- Nicoletta, V., Chavardès, R.D., Abo El Ezz, A., Cotton-Gagnon, A., Bélanger, V., and Boucher, J. 2023. Firelossrate: an R package for computing wildfire impacts on structures at the Wildland Urban interface. Available from SSRN 4391692.
- Noovo Info. 2023. Les pilotes d'avion-citerne se rapprochent de la limite de temps de vol selon la loi. Available from <https://www.noovo.info/nouvelle/les-pilotes-davion-citerne-bient-ot-a-la-limite-selon-la-loi.html> [accessed 21 December 2023].
- NRCan. 2020. Overview of Canada's forest industry. Available from <https://www.nrcan.gc.ca/our-natural-resources/forests-forestry/forests-industry-trade/overview-canadas-forest-industry/13311> [accessed 22 October 2023].
- NRCan. 2023. Statistical data. Available from <https://cfs.nrcan.gc.ca/statprofile/economicimpact/CA> [accessed 4 December 2023].
- Observatoire régional de recherche sur la forêt boréale [ORRFB]. 2023. Actes du Sommet sur les feux de forêt, 6–7 Novembre 2023. Université du Québec à Chicoutimi. 12.
- O'Neill, B., Oppenheimer, M., Warren, R., Hallegatte, S., Kopp, R.E., Pörtner, H.O., et al. 2017. “IPCC reasons for concern regarding climate change risks”. *Nat. Clim. Change*, **7**: 28–37. doi:10.1038/nclimate3179.
- Parisien, M.-A., Dawe, D.A., Miller, C., Stockdale, C.A., and Armitage, O.B. 2019. Applications of simulation-based burn probability modelling: a review. *Int. J. Wildl. Fire*, **28**: 913–926. doi:10.1071/WF19069.
- Parisien, M.A., Barber, Q.E., Bourbonnais, M.L., Daniels, L.D., Flannigan, M.D., Gray, R.W., et al. 2023. Abrupt, climate-induced increase in wildfires in British Columbia since the mid-2000s. *Commun. Earth Environ.* **4**: 309. doi:10.1038/s43247-023-00977-1.
- Parlee, B.L., Geertsema, K., and Willier, A. 2012. Social-ecological thresholds in a changing boreal landscape: insights from Cree knowledge of the Lesser Slave Lake region of Alberta. *Ecol. Soc.* **17**(2): 20. doi:10.5751/ES-04410-170220.
- Pau, M., Gauthier, S., Chavardès, R.D., Girardin, M.P., Marchand, W., and Bergeron, Y. 2022. Site index as a predictor of the effect of climate warming on boreal tree growth glob. *Global Change Biol.* **28**: 1903–1918. doi:10.1111/gcb.16030.
- Pau, M., Gauthier, S., Boulanger, Y., Ouzennou, H., Girardin, M.P., and Bergeron, Y. 2023. Response of forest productivity to changes in growth and fire regime due to climate change. *Can. J. For. Res.* **53**: 663–676. doi:10.1139/cjfr-2022-0207.
- Perrault-Hébert, M., Boucher, Y., Fournier, R., Girard, F., Auger, I., Thiffault, N., and Grenon, F. 2017. Ecological drivers of post-fire regeneration in a recently managed boreal forest landscape of eastern Canada. *Forest Ecol. Manage.* **399**: 74–81. doi:10.1016/j.foreco.2017.05.026.
- Public Safety Canada. 2023. The Canadian Disaster Database. Available from <https://www.publicsafety.gc.ca/cnt/rsrscs/cndn-dsstr-dtbs/index-en.aspx> [accessed 15 December 2023].
- Purdon, M., Brais, S., and Bergeron, Y. 2004. Initial response of understorey vegetation to fire severity and salvage-logging in the southern boreal forest of Québec. *Appl. Veg. Sci.* **7**: 49–60. doi:10.1111/j.1654-109X.2004.tb00595.x.
- Radio-Canada. 2023. Avions-citernes: des effectifs disparates et vieillissants pour affronter les feux. Available from <https://ici.radio-canada.ca/nouvelle/2006075/bombardiers-eau-essentiel-s-disparates-vieillissants> [accessed 21 December 2023].
- Raulier, F., Dhital, N., Racine, P., Tittler, R., and Fall, A. 2014. Increasing resilience of timber supply: how a variable buffer stock of timber can efficiently reduce exposure to shortfalls caused by wildfires. *For. Pol. Econ.* **46**: 47–55. doi:10.1016/j.forpol.2014.06.007.
- Remy, C.C., Hély, C., Blarquez, O., Magnan, G., Bergeron, Y., Lavoie, M., and Ali, A.A. 2017. Different regional climatic drivers of Holocene large wildfires in boreal forests of northeastern America. *Environ. Res. Lett.* **12**(3): 035005. doi:10.1088/1748-9326/aa5aff.
- Robinne, F.-N., Bladon, K.D., Miller, C., Parisien, M.-A., Mathieu, J., and Flannigan, M.D. 2018. A spatial evaluation of global wildfire-water risks to human and natural systems. *Sci. Total Environ.* **610–611**: 1193–1206. doi:10.1016/j.scitotenv.2017.08.112. PMID: 28851140.
- Rudolph, T.D., and Laidly, P.R. 1990. *Pinus banksiana* (Lamb.), Jack Pine. *In* *Silvics of North America*. Vol. 1. Edited by R.M. Burns and B.H. Honkala. Conifers, U.S.D.A. For. Serv. Agric. Handbk. 654. Washington, D.C. pp. 280–293.
- Rudolph, T.D., Drapeau, P., Imbeau, L., Brodeur, V., Légaré, S., and St-Laurent, M.-H. 2017. Demographic responses of boreal caribou to cumulative disturbances highlight elasticity of range-specific tolerance thresholds. *Biodivers. Conserv.* **26**: 1179–1198. doi:10.1007/s10531-017-1292-1.
- Saint-Germain, M., and Green, D.F. 2009. Salvage logging in the boreal and cordillera forests of Canada: integrating industrial and ecological concerns in management plans. *Forestry Chron.* **85**: 120–134. doi:10.5558/tfc85120-1.
- Saucier, J.-P., Gosselin, J., Morneau, C., and Grondin, P. 2010. Utilisation de la classification de la végétation dans l'aménagement forestier au Québec. *Rev. For. Française*, **62**(3–4): 428–438. doi:10.4267/2042/38956.
- Savage, D.W., Martell, D.L., and Wotton, B.M. 2010. Evaluation of two risk mitigation strategies for dealing with fire-related uncertainty in timber supply modelling. *Can. J. For. Res.* **40**(6): 1136–1154. doi:10.1139/X10-065.
- Sénécal, P., and Égré, D. 1999. Human impacts of the La Grande hydroelectric complex on Cree communities in Québec. *Impact Assess. Project Appraisal*, **17**(4): 319–329. doi:10.3152/147154699781767648.
- Shakeri, Z. 2024. Étude de la mosaïque forestière structurée par les feux dans la taïga de l'est du Canada. Ph. D. thesis, UQAR.
- Société de protection des forêts contre le feu [SOPFEU]. 2023a. Société de protection des forêts contre le feu. Available from sopfeu.qc.ca [accessed 15 December 2023].
- Société de protection des forêts contre le feu [SOPFEU]. 2023b. Le gouvernement du Québec annonce un soutien de 16 M\$ pour la SOPFEU. Available from <https://sopfeu.qc.ca/communiqués/le-gouvernement-du-quebec-annonce-un-soutien-de-16-m-pour-la-sopfeu/> [accessed 21 December 2023].
- Splawinski, T.B., Cyr, D., Gauthier, S., Jetté, J.P., and Bergeron, Y. 2019. Analyzing risk of regeneration failure in the managed boreal forest of northwestern Québec. *Can. J. For. Res.* **49**(6): 680–691. doi:10.1139/cjfr-2018-0278.
- St-Laurent, M.-H., Boulanger, Y., Cyr, D., Manka, F., Drapeau, P., and Gauthier, S. 2022. Lowering the rate of timber harvesting to mitigate impacts of climate change on boreal caribou habitat quality in eastern Canada. *Sci. Total Environ.* **838**: 156244. doi:10.1016/j.scitotenv.2022.156244.
- Statistics Canada. 2023a. Census profile, 2021 census of population. Available from <https://www12.statcan.gc.ca/census-recensement/2021/dp-pd/prof/index.cfm?Lang=E> [accessed 21 December 2023].
- Statistics Canada. 2023b. Canada's demographic estimates for July 1, 2023: record-high population growth since 1957. Available from <https://www150.statcan.gc.ca/n1/daily-quotidien/230927/dq230927a-eng.htm> [accessed 19 December 2023].
- Terrier, A., Girardin, M.P., Périé, C., Legendre, P., and Bergeron, Y. 2013. Potential changes in forest composition could reduce impacts of climate change on boreal wildfires. *Ecol. Appl.* **23**: 21–35. doi:10.1890/12-0425.1.
- The New York Times. 2023. Wildfire smoke—smoky air disrupts life in the northeast. Available from <https://www.nytimes.com/live/2023/06/07/us/canada-wildfires-air-quality-smoke> [accessed 4 December 2023].

- Thorn, S., Bässler, C., Brandl, R., Burton, P.J., Cahall, R., Campbell, J.L., et al. 2018. Impacts of salvage logging on biodiversity: a meta-analysis. *J. Appl. Ecol.* **55**(1): 279–289. doi:[10.1111/1365-2664.12945](https://doi.org/10.1111/1365-2664.12945).
- Thorn, S., Chao, A., Georgiev, K.B., Müller, J., Bässler, C., Campbell, J.L., et al. 2020. Estimating retention benchmarks for salvage logging to protect biodiversity. *Nat. Commun.* **11**: 4762. doi:[10.1038/s41467-020-18612-4](https://doi.org/10.1038/s41467-020-18612-4).
- Tremblay, J.A., Boulanger, Y., Cyr, D., Taylor, A.R., Price, D.T., and St-Laurent, M.-H. 2018. Harvesting interacts with climate change to affect future habitat quality of a focal species in eastern Canada's boreal forest. *PLoS ONE*, **13**(2): e0191645. doi:[10.1371/journal.pone.0191645](https://doi.org/10.1371/journal.pone.0191645).
- Tymstra, C., Stocks, B.J., Cai, X., and Flannigan, M.D. 2020. Wildfire management in Canada: review, challenges and opportunities. *Progr. Disaster Sci.* **5**: 100045. doi:[10.1016/j.pdisas.2019.100045](https://doi.org/10.1016/j.pdisas.2019.100045).
- United Nations Office for Disaster Risk Reduction [UNDRR]. 2019. Global assessment report on disaster risk reduction, United Nations Office for Disaster Risk Reduction (UNDRR), Geneva.
- United Nations Office for Disaster Risk Reduction [UNDRR]. 2022. Global Assessment Report on Disaster Risk Reduction 2022: Our World at Risk: Transforming Governance for a Resilient Future. Geneva.
- Van Wagner, C.E. 1987. Development and structure of the Canadian Forest Fire Weather Index System. *Can. For. Serv., Ottawa, Ont. For. Tech. Rep.* 35. 37p.
- Venier, L.A., Thompson, I.D., Fleming, R., Malcolm, J., Aubin, I., Trofymow, J.A., et al. 2014. Effects of natural resource development on the terrestrial biodiversity of Canadian boreal forests. *Environ. Rev.* **22**(4): 457–490. doi:[10.1139/er-2013-0075](https://doi.org/10.1139/er-2013-0075).
- Vincent, L.A., Hartwell, M.M., and Wang, X.L. 2020. A third generation of homogenized temperature trend analysis and monitoring changes in Canada's climate. *Atmos. Ocean*, **58**: 173–191. doi:[10.1080/07055900.2020.1765728](https://doi.org/10.1080/07055900.2020.1765728).
- Walker, D.A., Forbes, B.C., Leibman, M.O., Epstein, H.E., Bhatt, U.S., Comiso, J.C., et al. 2011. Cumulative effects of rapid land-cover and land-use changes on the Yamal Peninsula. *In* Eurasian Arctic land cover and land use in a changing climate. Issue May 2014. Edited by G. Gutman and A. Reissel. Springer. pp. 207–236. doi:[10.1007/978-90-481-9118-5](https://doi.org/10.1007/978-90-481-9118-5).
- Wang, J., Taylor, A.R., and D'Orangeville, L. 2022. Warming induced tree growth may help offset increasing disturbance across the Canadian boreal forest. *Proc. Natl. Acad. Sci.* **120**: 1–10.
- Wazneh, H., Gachon, P., Laprise, R., De Vernal, A., and Tremblay, B. 2021. Atmospheric blocking events in the North Atlantic: trends and links to climate anomalies and teleconnections. *Clim. Dyn.* **56**: 2199–2221. doi:[10.1007/s00382-020-05583-x](https://doi.org/10.1007/s00382-020-05583-x).
- Williamson, T.B., Price, D.T., Beverly, J., Bothwell, P.M., Parkins, J.R., Patriquin, M.N., et al. 2007. A framework for assessing vulnerability of forest-based communities to climate change. Information Report NOR-X-414E. Natural Resources Canada, Canadian Forest Service, Northern Forestry Centre, Edmonton, Alberta. 50 p.
- Wotton, B.M., Flannigan, M.D., and Marshall, G.A. 2017. Potential climate change impacts on fire intensity and key wildfire suppression thresholds in Canada. *Environmental Research Letters*, **12**(9): 095003. doi:[10.1088/1748-9326/aa7e6e](https://doi.org/10.1088/1748-9326/aa7e6e).
- Zhao, F., and Liu, Y. 2019. Atmospheric circulation patterns associated with wildfires in the monsoon regions of China. *Geophys. Res. Lett.* **46**: 4873–4882. doi:[10.1029/2019GL081932](https://doi.org/10.1029/2019GL081932).