

1 Title

2 **GROWTH RESPONSE AND WATER RELATIONS OF THREE-YEAR-OLD**

3 **PLANTED BLACK SPRUCE AND JACK PINE SEEDLINGS IN SITE**

4 **PREPARED LICHEN WOODLANDS**

5

6 François Hébert^a, Jean-François Boucher^{b*}, Pierre Y. Bernier^c, Daniel Lord^{dt}

7

8 ^aUniversité Laval, Faculté de foresterie et géomatique, Département des
9 sciences du bois et de la forêt, Pavillon Abitibi-Price, Québec, Canada, G1K 7P4

10

11 ^bUniversité du Québec à Chicoutimi, Département des Sciences fondamentales,
12 555 boul. Université, Chicoutimi, Québec, Canada, G7H 2B1

13

14 ^cNatural Resources Canada, Canadian Forest Service, Laurentian Forestry
15 Centre, P.O. Box 3800, Quebec, Quebec, Canada G1V 4C7

16

17 *Corresponding author: Tel.: (418) 545-5011 ext. 5385. Fax.: (418) 545-5012.
18 Email: jean-francois_boucher@uqac.ca

19

20 †Reprint request: Tel.: (418) 545-5011 ext. 5064. Fax: (418) 545-5012. Email:
21 Daniel_Lord@uqac.ca

22

1 **Abstract**

2

3 Black spruce-lichen woodlands are low tree density stands within the
4 closed crown, North American boreal forest that represent a diverging post-fire
5 type of black spruce forest. As natural densification of lichen woodlands has
6 never been observed, plantation remains the only way of shifting these stands to
7 closed canopy stands. The objective of the study was to evaluate site preparation
8 effects on growth and water relations of black spruce (*Picea mariana* (Mill.)
9 B.S.P.) and jack pine (*Pinus banksiana* Lamb.) seedlings in black spruce-lichen
10 woodlands (LW), compared to managed black spruce-feathermoss stands
11 (BSFM). Site preparation treatments in LW were no preparation (LWno), patch
12 scarification (LWps), and disk scarification (LWds). The operationally managed
13 BSFM stands stood for the control. Results from the third growing season
14 indicate that soil water availability in intact or lightly prepared patch scarified LW
15 is a limiting variable for seedling growth for both black spruce and jack pine
16 seedlings. However, once LW are prepared with disk scarification, this planting
17 check factor is significantly reduced, to the point of being equal to BSFM in terms
18 of water availability and seedling water relations. The significant seedling growth
19 difference, in favor of BSFM, might be a consequence of the lower level of pre-
20 plantation disturbance in LW, compared to BSFM. Our results also suggest that
21 jack pine, through higher water stress tolerance, could constitute a wise
22 silvicultural choice over black spruce, especially with LW established on coarse
23 drought-prone material.

1

2 **Keywords:** *Picea mariana*; *Pinus banksiana*; Lichen woodland; Site preparation;
3 Water relations; Afforestation

1 **1. Introduction**

2

3 Black spruce-lichen woodlands (hereafter shortened to lichen woodlands)

4 are low tree density stands, scattered within the closed-crown black spruce-

5 feathermoss forest sub-zone of the Québec's boreal forest zone (Payette, 1992;

6 Bergeron, 1996). These open stands are characterized by a sparse tree layer of

7 black spruce (*Picea mariana* (Mill.) B.S.P.), and a lichen mat (*Cladina* spp.) often

8 mixed with ericaceous shrub species (mainly *Kalmia angustifolia* L., *Ledum*

9 *groenlandicum* Oeder, and *Vaccinium angustifolium* Ait.). Lichen woodlands

10 apparently result from repeated catastrophic disturbances (wildfire and insect

11 defoliation) in formerly closed-crown black spruce-feathermoss (BSFM) stands

12 within the boreal forest, and contribute to the progressive and irreversible loss of

13 BSMF stands (Payette, 1992; Payette et al., 2000; Gagnon and Morin, 2001).

14 Presently, lichen woodlands make up to 10% of total forested lands within the

15 closed-crown black spruce-feathermoss forest sub-zone, based on ecoforestry

16 maps of the Québec's Ministry of Natural resources and wildlife (MRNF)

17 (Anonymous, 2005). There is no evidence of natural redensification of lichen

18 woodlands, i.e. a shifting to a closed-crown black spruce-feathermoss stand

19 (Payette, 1992; Riverin and Gagnon, 1996).

20

21 Lichen woodlands have not been studied yet for their afforestation

22 potential probably because of their low timber yield (Riverin and Gagnon, 1996;

23 Gagnon and Morin, 2001). Limitations to afforestation have been identified in

1 *Kalmia-Ledum* woodlands of the North American boreal forest, which share some
2 similarities with lichen woodlands because of their low tree density and the
3 presence of ericaceous shrubs (Mallik, 1993; Inderjit and Mallik, 1996; Yamasaki
4 et al., 1998). Reduced soil fertility, allelopathic interference and impacts on
5 mycorrhizal colonization have been identified as potential limitations to young
6 spruce growth and survival in *Kalmia-Ledum* woodlands (Zhu and Mallik, 1994;
7 Inderjit and Mallik, 1996; 2002; Parker et al., 1997; Yamasaki et al., 1998; 2002;
8 Thiffault et al., 2003). By contrast, influence of lichens on conifer seedlings have
9 been rarely studied, and mechanisms of interference have not been yet found, if
10 any exist (Kershaw and Rouse, 1971; Sirois, 1993; Steijlen et al., 1995). Water
11 stress is major factor that can contribute to growth check in young conifer
12 plantations (Burdett et al., 1984; Bernier, 1993). Considering that lichen
13 woodlands are reputed drought prone habitats, early afforestation efforts in these
14 environments should also include some investigation on water relations of
15 planted conifers (Mallik, 1991; 1993; Prévost, 1992; Bergeron, 1996).

16

17 Silvicultural approaches, such as site preparation, can potentially
18 overcome water stress induced growth check in planted seedlings. For instance,
19 disk scarification, the most common site preparation treatment in Québec, can
20 improve short term growth conditions by diminishing competing vegetation
21 density, and thus, improving water availability for seedling roots (Bassman, 1989;
22 Fleming et al., 1994; Brais 2001). By exposing the mineral horizon of soil and
23 increasing soil temperature, disk scarification is also known to improve water

1 absorption through increased root growth, root cell permeability and hydraulic
2 conductivity, and decreased soil water viscosity (Grossnickle and Heikurinen,
3 1989; Bowen, 1991; Boucher et al., 1998 and 2001). Scarification can also have
4 negative impacts, principally through increased soil surface evaporation and,
5 consequently, water deficits in small seedlings (Stathers and Spittlehouse, 1990;
6 Prévost, 1992). Patch scarification represents an alternate site preparation
7 method on those sites where disk preparation is impractical, with likely similar,
8 but little studied impacts on soil conditions or on seedling water relations
9 (Örlander, 1986; Tremblay, 1996; Mäkitalo and Hyvönen, 2004). Use of a more
10 water stress tolerant conifer like jack pine (*Pinus banksiana* Lamb.), instead of
11 less tolerant black spruce, is an additional silvicultural choice that may improve
12 success on drought prone sites (Grossnickle and Blake, 1986).

13

14 This text presents the first results obtained from an experimental
15 plantation network established in 1999 throughout the BSFM sub-zone and
16 designed to test the efficiency of various site preparation methods in the
17 promotion of survival and growth of planted seedlings in lichen woodlands. The
18 objectives of the study were 1) to determine if site preparation of lichen
19 woodlands could generate levels of survival and growth of planted seedlings at
20 least comparable to those found in seedlings planted in harvested black spruce-
21 feathermoss stands, 2) to determine if there was a gain to be made by using the
22 more drought-tolerant jack pine seedlings instead of the black spruce seedlings
23 in the regeneration of lichen woodlands, and 3) to determine if differences in

1 survival and growth found between treatments could be related to water stress.

2

1 **2. Materials and methods**

2

3 *2.1. Study site description*

4

5 The experiment was carried out in five experimental blocks split over two
6 sites within black spruce-feathermoss forest of the Quebec's boreal forest
7 (Bergeron, 1996). Both sites were located in the central part of the black spruce-
8 feathermoss forest, one site near the west shore of the Mistassibi River (centered
9 at 50°04'49" N, 71°59'44" W), and the other site near the east shore of the
10 Péribonka Lake (centered at 50°10'32" N, 71°09'49"W). Site selection was based
11 on two main criteria: i) proximity of black spruce-feathermoss and lichen
12 woodland stands of comparable slope, aspect and topography, and ii) nearby
13 planned harvesting and site preparation operations.

14

15 Each of the five blocks included BSFM and LW stands. In all blocks,
16 BSFM stands were originally of about 120 years of age, and were dominated by
17 mature black spruce (*Picea mariana* (Mill.) B.S.P.), with a small component of
18 jack pine (*Pinus banksiana* Lamb.). Pre-harvesting crown cover – expressed as
19 the ground area covered by the crowns of trees as delimited by the vertical
20 projection of crown perimeters – of BSFM stands was between 60 and 80% in
21 blocks 1 and 2, between 40 and 60% in blocks 3 and 4, and between 25 to 40%
22 in block 5. The understory included black spruce advance regeneration, a sparse
23 shrub layer of *Kalmia angustifolia* L., *Ledum groenlandicum* Oeder, *Vaccinium*

1 *myrtilloides* Michx., *Vaccinium angustifolium* Ait., and *Gaultheria hispida* (L.)
2 Mühl. ex Bigel., and a dense mat of mosses, mostly *Pleurozium schreberi* (Brid.)
3 Mitt., *Hylocomium splendens* (Hedw.) B.S.G., *Ptilium crista-castrensis* (Hedw.)
4 De Not., *Rhytidiodelphus triquetrus* (Hedw.) Warnst., *Polytrichum* spp., with
5 some *Sphagnum* spp.

6

7 By contrast, the adjoining lichen woodlands were characterized in all
8 blocks by a tree cover of black spruce of less than 25% and a ground cover of
9 lichen greater than 40% and dominated by *Cladina mitis* (Sandst.) Hustich,
10 *Cladina stellaris* (Opiz) Brodo, and *Cladina rangiferina* (L.) Nyl. The lichen
11 woodlands also had a dense shrub layer composed of the same species found in
12 the BSFM stands.

13

14 All experimental stands were on moderately deep (50-100 cm) to deep (>
15 100 cm) coarse glacial till deposits, overtopped by a mor humus with humo-ferric
16 podzolic profiles, except for two lichen woodlands in one block of the Mistassibi
17 site which were on deep glaciofluvial outwashes. Results presented were most
18 likely not influenced by this difference in deposit types between plots of this
19 block, since soil texture and drainage were similar (results not shown). Drainage
20 on both sites was good to moderate. See also Saucier et al. (1994) and Bergeron
21 (1996) for a general description of the black spruce-feathermoss forest in
22 Quebec's boreal forest zone).

23

1 Climate in this central part of the spruce-feathermoss forest is cool
2 continental, with a mean annual temperature of 0.0 °C (± 1.3 °C) and a total
3 precipitation of 961 mm, with 302 mm (31 %) as snow (1971-2000 data from the
4 Chapais weather station, 49° 47' N – 74° 51' O) (Environment Canada, 2004).
5 Data loggers set up 40 km West of the Mistassibi River site show that climate
6 was generally close to normal during the course of the experimental period
7 (1999-2002), with average temperature of 0.6 °C (± 0.9 °C) and total rainfall of
8 603 mm.

9

10 *2.2. Experimental design and treatments*

11

12 The experimental design was a five-block, 4 X 2 split-plot design, with four
13 different site preparation/stand combinations (hereafter “treatment(s)”) and two
14 planted species within each treatment. The four treatments included three types
15 of site preparation in lichen woodlands (LW) and one operationally managed
16 BSFM stand, which included clearcut harvesting, followed by disk scarification
17 (hydraulic TTS) both in and between the skid trails. Because of the combination
18 of disk scarification and harvesting operations, approximately 30 % of the BSFM
19 stand area was disturbed twice (Harvey and Brais, 2002). Harvesting/scarification
20 operations were scheduled as follows: block 1 in 1997/1997, blocks 2 and 3 in
21 1994/1998, and blocks 4 and 5 in 1999/1999. Notice that only BSFM stands were
22 harvested before performing treatments, so that these experimental plots in
23 BSFM stands should be interpreted as the point of reference in silvicultural terms.

1 The site preparations in LW were disk scarification (LWds), patch
2 scarification (LWps) and no preparation (LWno). Disk scarification was carried
3 out by a hydraulic TTS disk trencher on the Mistassibi River site, and by a non-
4 hydraulic TTS disk trencher on the Pérignon Lake site. The furrows were 2 m
5 apart and had a mean width of 67 cm and a mean depth of 18 cm (n=24). As a
6 result of disk scarification, approximately 34% of total stand area was disturbed.
7 Patch scarification was carried out in 1999 using a cutter-type portable scarifier
8 that removed the aboveground portion of the brush and reduced the humus layer
9 thickness. The 15 cm-radius patches were done on a 2x2 m spacing, so that
10 approximately 2 % of total stand area was disturbed.

11

12 Containerized black spruce (*Picea mariana* (Mill.) B.S.P.) and jack pine
13 (*Pinus banksiana* Lamb.) seedlings were produced from local seeds using 67-
14 cavity-containers with 50 cm³ root plug volume per cavity (IPL inc., Saint-Damien,
15 Qc, Canada). Mean total height and stem diameter at planting time were of 174
16 mm and 2.4 mm, and of 168 mm and 2.0 mm for jack pine and black spruce
17 respectively (MRNF, unpublished data). Approximately 200 seedlings of each
18 species were planted at a 2x2 m spacing into two randomized subplots within
19 each one of the main treatment plot. Planting took place in late summer-early fall
20 of 1999. Seedlings were planted at the sides of the disk scarification trails, in the
21 middle patches of the scarification patches, and directly into the forest
22 floor/humus interface of the no preparation plots.

23

1 2.3. *Soil and seedling measurements*

2

3 Physiological measurements were taken on one randomly chosen
4 seedling per subplot for each of the following four sampling periods of the 2002
5 active growth season: June 10-13, July 24-26, August 8-13 and September 17-
6 18. Measurements were done on clear days preceded by at least one day without
7 rain. On each sampling date, predawn water potential (Ψ_{pd}) and shoot
8 conductance to water vapor (g_{wv}) were measured on the top 10 cm of the main
9 shoot excised from each selected seedling. No measurement of Ψ_{pd} could be
10 taken during the August sampling period at the Péribonka site, and during the
11 September sampling period on both sites. Sites were visited on consecutive
12 days, the distance between them preventing same-day measurements.

13

14 For Ψ_{pd} measurements, each excised shoot was rapidly put in a plastic
15 bag and placed in a cooler with ice. Measurements were made using a pressure
16 chamber (PMS Instruments, Corvallis, OR) (Scholander et al., 1965). All shoots
17 in a block were collected within 45 minutes and one complete site was collected
18 and measured approximately two to three hours before dawn.

19

20 For measurements of g_{wv} , each excised shoot was placed in a sealed
21 plastic bag containing a wet sponge, and was left exposed to natural light before
22 measurement. Measurements were taken using a LI-6200 portable
23 photosynthesis system (LI-COR, Inc., Lincoln, NE) with a 0.25 l cuvette, and an

1 halogen lamp approx. 15 cm over the cuvette when the sunlight was not
2 providing saturation ($\geq 1200 \mu\text{mol photons m}^{-2} \text{s}^{-1}$) due to passing clouds. Shoots
3 were measured in a central location less than 30 minutes after excision and all
4 shoots within a block were collected within 45 minutes. Preliminary tests showed
5 that excised shoots from black spruce and jack pine seedlings can be stored in
6 this manner up to 45 minutes without any change in g_{wv} (data not shown). All
7 measurements within a site were taken between 10h00 and 12h00 E.T.

8

9 For each seedling used for g_{wv} measurements, a sub-sample of needles
10 placed in the cuvette was used to determine total foliar surface area. Needle
11 surface area estimation method was similar to that in Bernier et al. (2001) with
12 balsam fir (*Abies balsamea* (L.) Mill.). The ratio of estimated area (ratio of
13 estimated area from the needle central section / estimated area from several
14 needle sections) obtained for black spruce was 0.892 and 0.915 for jack pine and
15 was used to estimate total surface area of the sub-samples. The needle sub-
16 samples used for surface area estimation were then dried at 65 °C for 48 hours
17 to obtain leaf dry mass per unit leaf area (LMA). The LMA calculated was then
18 used to find total foliar area in the cuvette; each g_{wv} measurement was then
19 recalculated with actual total needle surface area.

20

21 The remaining part of each seedling used for both g_{wv} and Ψ_{pd}
22 measurements was dug out for morphological measurements in the lab. After
23 having carefully washed the root system of each seedling, total seedling height

1 (H_t; nearest mm), stem hypocotyl diameter (D_s; nearest one-tenth mm), root,
2 stem and foliar dry mass (65° C for 48 hours) were measured. Total seedling dry
3 mass (DM_t), foliar to shoot dry mass ratio (FS_{dm}), and root to shoot dry mass ratio
4 (RS_{dm}) were calculated from these dry mass measurements. Survival of
5 seedlings within each subplot was determined at the end of the growing season
6 from one hundred of the initial seedlings within each subplot.

7

8 For each selected seedling within each subplot at each of the four
9 sampling date throughout the 2002 growing season (June 10-13, July 24-26,
10 August 8-13 and September 17-18), three 50 ml samples were collected, using a
11 soil sampler (Soil Moisture Inc., Santa Barbara, CA), within a radius of 15 cm of
12 each seedling at root plug depth. The samples were either from organic and/or
13 mineral horizons, depending on seedlings' respective root plug environment, and
14 were bulked for each seedling (at each sampling date) and used to obtain soil
15 water content (SWC) by the gravimetric method (Rundel and Jarrell, 1991) and
16 organic matter percentage (%OM) by the loss-on-ignition method (Nelson and
17 Sommers, 1983) needed to characterize the microsite of each planted seedling.

18

19 *2.4. Statistical analyses*

20

21 Analysis of variance (ANOVA) on incomplete split-split-plot design was
22 used for physiological and soil variables, with site treatments at the main plot
23 level, planted tree species at the subplot level and sampling dates at the sub-

1 subplot level. An incomplete split-plot design was used for morphological
2 variables, ratios, and survival, using the September data set (no sampling date
3 factor). The incomplete block structure was due to missing black spruce
4 seedlings in LWno plot of two blocks, and to a removed BSFM plot of one block,
5 owing to its particular stand history (wildfire in 1996 before harvesting).

6

7 Relative growth rate (RGR_{dm}) was used to analyze treatment or species
8 effects on seedling growth, using the Poorter and Lewis method (1986), based on
9 analysis of variance, with ln-transformed total dry mass as the dependent
10 variable. A significant Treatment *Time interaction indicates a difference in
11 relative growth rates between treatments. Calculated ratios (FS_{dm} , RS_{dm} , and
12 LMA) were adjusted by the Bauce et al. method (1994) in order to respect
13 fundamental statements of ANOVA (Zar 1999). This method standardizes the
14 ratios by adjusting the numerators of each ratio to a common denominator before
15 conducting the ANOVA's. Homogeneity of variance was verified for all data by
16 visual analysis of residues (Devore and Peck, 1994), and logarithmic
17 transformations were performed when necessary to homogenize the variance.
18 Seedling survival was transformed using $\text{Sin}^{-1}(\sqrt{(\text{survival} \cdot .01) \cdot (180 \cdot \pi^{-1})})$.

19

20 ANOVA's were performed using the MIXED procedure of SAS 8.1
21 software (SAS Institute, Cary, NC). The SLICE command in the MIXED
22 procedure was made after the ANOVA when the Treatment*Species interaction
23 was significant, in order to determine if each of both species was influenced

1 significantly by the treatments. This procedure allows one to make a separate
2 ANOVA for each species to determine if there were differences among
3 treatments. Differences among treatments were determined by three a-priori
4 contrasts (Steel and Torrie, 1980) aimed at, first, evaluating the respective impact
5 of both site preparations tested within LW and, secondly, verifying how disk
6 scarified LW compares with the point of reference, BSFM stands, since LWdisk
7 corresponds to the nearest silvicultural approach compared to operationally
8 managed BSFM: i) LWps vs LWno; ii) LWds vs LWno; iii) LWds vs BSFM.
9 Differences were considered significant at $P < 0.05$.

10

1 **3. Results**

2

3 *3.1. Morphological variables and survival*

4

5 In lichen woodlands (LW), patch scarification did not influence seedling
6 growth, but disk scarification significantly increased growth of both species
7 compared to the no site preparation (LWno) plots (Table 1 and Fig. 1). Values of
8 total height (H_t), stem hypocotyl diameter (D_s), and total dry mass (DM_t) were,
9 respectively, 27%, 63% and 200% greater in the LWds plot than in the LWno
10 plots. However, seedling growth variables in LWds were not as high as those
11 found for the seedlings in the black spruce-feathermoss stands (BSFM). Values
12 of total height (H_t), stem hypocotyl diameter (D_s), and total dry mass (DM_t) were,
13 respectively, 18%, 29%, and 52% higher in BSFM than in LWds (Table 1 and Fig.
14 1 a,c,e). Growth of both species followed a similar pattern (no significant
15 treatment*species interaction), but jack pine seedlings showed overall higher
16 growth values than black spruce seedlings, with H_t , D_s , and DM_t respectively
17 37%, 47%, and 35% higher in jack pine than in black spruce (Table 1, Fig. 1 b, d,
18 f). Biomass allocation in seedlings was not influenced by the different site
19 preparations in LW (Table 1), but allocation to root (root-to-shoot dry mass ratio
20 or RS_{dm}) of seedlings was significantly higher in LWds plots than in BSFM plots
21 (Table 2, Fig. 1g). Neither the above-ground allocation to foliage (FS_{dm}) nor the
22 leaf mass per unit area (LMA) were influenced by treatments (Table 2). No
23 interaction between time and treatment or species was found for the In

1 transformed dry masses indicating that relative growth rate (RGR_{dm}) was not
2 influenced by either treatment or species (Table 3).

3

4 Both patch and disk site preparations in LW significantly increased the
5 survival of seedlings after three years of growth in the field, regardless of the tree
6 species (Table 1, Fig. 1h). Seedling survival in LWds was equal to that of
7 seedlings in BSFM (Table 1, Fig. 1h).

8

9 **Table 1:** Summary of ANOVA (F and P -values) for total height (H_t), shoot diameter (D_s), total dry
10 mass (DM_t) and survival in black spruce and jack pine seedlings after three growing seasons.
11 Bold indicates significance ($P < 0.05$). ndf = numerator degrees of freedom, LW= lichen
12 woodlands (subscripts ps for patch scarification, ds for disk scarification and no for no
13 preparation), and BSFM = black spruce-feathermoss stands.

Source of variation	H_t			D_s			DM_t			Survival [†]	
	ndf	F	P	F	P	F	P	F	P	F	P
Treatment (T)	3	33.59	< 0.0001	38.20	< 0.0001	36.91	< 0.0001	12.71			0.0004
Species (S)	1	36.44	0.0006	43.93	< 0.0001	12.47	0.0008	1.31			0.2972
T*S	3	0.53	0.6658	2.64	0.0573	0.56	0.6431	1.88			0.1907
<i>Contrasts</i>											
LWps vs LWno	1	3.82	0.0555	0.08	0.7841	0.48	0.4901	11.38			0.0050
LWds vs LWno	1	35.07	< 0.0001	28.73	< 0.0001	45.33	< 0.0001	35.36			< 0.0001
LWds vs BSFM	1	13.44	0.0005	15.44	0.0002	0.48	0.0514	1.75			0.2081

15 *Ln transformed data

16 †Transformed data (($\text{Sin}^{-1}(\sqrt{(\text{survival}/100))} \times 180/\pi$)

17

18

1

2 **Table 2:** Summary of ANOVA (*F* and *P*-values) for root to shoot ratio (RS_{dm}), foliar dry mass to
 3 shoot dry mass ratio (FS_{dm}) and leaf mass per unit leaf area (LMA) in black spruce and jack pine
 4 seedlings during the September sampling period. Bold indicates significance (*P* < 0.05).
 5 Abbreviations are as in Table 1.

6

Source of variation	RS _{dm}			FS _{dm}			LMA	
	ndf	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	
Treatment (T)	3	2.99	0.0477	0.51	0.6746	1.68	0.1933	
Species (S)	1	0.95	0.3386	0.74	0.3934	1.75	0.1970	
T*S	3	2.15	0.1168	0.57	0.6343	0.16	0.9226	
<i>Contrasts</i>								
LWps vs LWno	1	0.00	0.9515					
LWds vs LWno	1	1.70	0.2031					
LWds vs BSFM	1	8.97	0.0057					

7

8 **Table 3:** Summary of ANOVA (*F* and *P* values) for total dry mass relative growth rate (RGR_{dm}) in
 9 black spruce and jack pine seedlings throughout the third growing season. Bold indicates
 10 significance (*P* < 0.05). Abbreviations are as in Table 1.

11

Source of variation	RGR _{dm}		
	ndf	<i>F</i>	<i>P</i>
Treatment (T)	3	27.85	< 0.0001
Species (S)	1	5.21	0.0725
T*S	3	1.90	0.1299
Date (D)	3	9.00	< 0.0001
D*T	9	1.34	0.2189
D*S	3	2.02	0.1117
T*S*D	9	0.58	0.8120

12

13

14




Fig. 1. Significant treatment and species effects on total height (a and b), stem diameter (c and d), total dry mass (e and f), root to shoot ratio (g), and survival (h) from the September sampling period. Each bar in (a), (c), (e), (g), and (h) represents the mean from 5 blocks, 2 species, and 2 samples per plot ($n=20$ for LWds and LWps, and $n=16$ for LWno and BSFM), and from 5 blocks, 4 treatments, and 2 samples per plot in (b), (d), and (f), ($n=34$ for BS, and $n=38$ for JP). Horizontal lines above the bars show significant treatment differences revealed by the contrasts (see Tables 1 and 2). Abbreviations are as in Table 1, plus BS = black spruce, JP = jack pine.


1
2
3
4
5
6
7
8
9

1 3.2. *Physiological variables*
 2
 3 Shoot predawn water potential (Ψ_{pd}) in seedlings was significantly
 4 increased by disk scarification in LW throughout the season, with values of Ψ_{pd} in
 5 LWds plots 47 % higher than that in LWno plots (Table 4, Fig. 2a). Patch
 6 scarification in LW did not result in higher values of Ψ_{pd} , compared to that in
 7 LWno, and values measured in seedlings in LWds plots were equal to that in
 8 BSFM stands (Table 4, Fig. 2a). Values of Ψ_{pd} in jack pine were 21 % higher than
 9 those in black spruce seedlings (Table 4, Fig 2b). Treatments had no effect on
 10 stomatal conductance for water vapor (g_{wv}) in black spruce seedlings ($0.0395 \pm$
 11 $0.0029 \text{ mol m}^{-2} \text{ s}^{-1}$). However, g_{wv} of jack pine seedlings was significantly greater
 12 in LWds than in LWno (Table 4, Fig. 3).

13
 14 **Table 4:** Summary of ANOVA (F and P -values) for predawn water potential (Ψ_{pd}) and stomatal
 15 conductance for water vapor (g_{wv}) measured in black spruce and jack pine seedlings during four
 16 sampling periods of the third growing season. Bold indicates significance ($P < 0.05$).
 17 Abbreviations are as in Table 1.
 18


Source of variation	Ψ_{pd}			g_{wv}		
	ndf	F	P	ndf	F	P
Treatment (T)	3	5.58	0.0030	3	0.75	0.5288
Species (S)	1	6.16	0.0182	1	1.83	0.1818
T*S	3	1.59	0.2101	3	2.84	0.0462
Date (D)	2	9.18	0.0006	3	5.95	0.0014
D*T	6	0.32	0.9234	9	0.63	0.7638
D*S	2	2.72	0.0803	3	1.45	0.2381
T*S*D	6	0.53	0.7852	9	0.90	0.5350
<i>Contrasts*</i>						
LWps vs LWno	1	0.07	0.8000	1	0.89	0.3501
LWds vs LWno	1	8.82	0.0052	1	10.00	0.0029
LWds vs BSFM	1	0.05	0.8236	1	0.13	0.7191

19 *Contrasts on g_{wv} were after a separate ANOVA made only for jack pine.
 20



1

2 **Fig. 2.** Significant treatment (a) and species (b) effects on seedling predawn water potential
 3 measured throughout the 2002 growing season. Each bar represents the mean from 5 blocks, 2
 4 species, and 3 dates in (a) ($n=26$ for LWds and LWps, $n=22$ for LWno, and $n=20$ for BSFM), and
 5 from 5 blocks, 4 treatments, and 3 dates in (b) ($n=45$ for BS, and $n=49$ for JP). Horizontal lines
 6 above the bars show significant treatment differences revealed by the contrasts (see Table 4).
 7 Abbreviations are as in Table 1, plus BS = black spruce, JP = jack pine.



8

9 **Fig. 3.** Significant interaction between treatments and species on seedling stomatal conductance
 10 for water vapor measured throughout the 2002 growing season. Each bar represents the mean
 11 from 5 blocks and 4 dates (for BS $n=20$ for LWds and LWps, $n=16$ for BSFM, and $n=12$ for LWno;
 12 for JP $n=20$ for LWds, LWps, and LWno, and $n=16$ for BSFM). Horizontal lines above the bars
 13 show significant treatment differences revealed by the contrasts (see Table 4). Abbreviations are
 14 as in Table 1, plus BS = black spruce, JP = jack pine.

15

16

1 3.3. Soil variables

2

3 Patch scarification did not influence the gravimetric soil water content
4 (SWC) in LW plots compared to LWno, but disk scarification significantly
5 increased it throughout the season (Table 5, Fig. 4). All three factors significantly
6 interacted to control organic matter percentage (%OM) in seedling soil
7 microsites. Organic matter percentage (%OM) was highly variable and unstable
8 across all treatments, especially in BSFM stands (Table 5, Fig. 5). Overall, patch
9 scarified sites had higher %OM than disk-scarified sites (Fig. 5).

10

11 **Table 5:** Summary of ANOVA (*F* and *P*-values) for gravimetric soil water content (SWC) and
12 percentage of soil organic matter (%OM) measured during four sampling periods of the third
13 growing season. Bold indicates significance (*P* < 0.05). Abbreviations are as in Table 1.

Source of variation	ndf	SWC		%OM	
		<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>
Treatment (T)	3	3.09	0.0418	1.17	0.3361
Species (S)	1	1.95	0.1726	1.26	0.2714
T*S	3	0.37	0.7761	0.13	0.9441
Date (D)	3	1.13	0.3724	0.61	0.6102
D*T	9	0.85	0.5703	0.81	0.6134
D*S	3	0.51	0.6779	1.64	0.1830
T*S*D	9	0.54	0.8383	2.56	0.0092
<i>Contrasts</i>					
LWps vs LWno	1	0.05	0.8266		
LWds vs LWno	1	5.72	0.0238		
LWds vs BSFM	1	2.99	0.1319		

15




Fig. 4. Significant treatment effects on soil water content measured throughout the 2002 growing season. Each bar represents the mean from 5 blocks, 4 dates, 2 species, and 2 samples per plot ($n=80$ for LWds and LWps, and $n=64$ for BSFM and LWno). Horizontal lines above the bars show significant treatment differences revealed by the contrasts (see Table 5). Abbreviations are as in Table 1, plus BS = black spruce, JP = jack pine.




Fig 5. Significant interaction between treatments, species microsite, and dates on soil organic matter content measured throughout the 2002 growing season. Each bar represents the mean from 5 blocks, and 2 samples per plot (for BS in each date, $n=10$ for LWds and LWps, $n=8$ for BSFM, and $n=6$ for LWno; for JP in each date, $n=10$ for LWds, LWps, and LWno, and $n=8$ for BSFM). Abbreviations are as in Table 1, plus BS = black spruce, JP = jack pine.

1 **4. Discussion**

2

3 *4.1. Site preparation impacts in lichen woodlands*

4

5 Disk scarification and patch scarification produce contrasting results with
6 respect to the soil environment, as well as the physiological response of planted
7 conifer seedlings in lichen woodlands (LW). Disk scarification appears to have a
8 greater ability to improve microsite water availability, resulting in higher seedling
9 growth, survival, and water potential, for both black spruce and jack pine. Patch
10 scarification, on the contrary, appears to provide no incremental gains in growth
11 compared to unscarified LW sites, in spite of its ability to improve seedling
12 survival. The beneficial impact of disk scarification on both growth and survival of
13 young planted conifers, especially during the establishment phase, has been
14 shown in a number of studies (Bassman, 1989; Grossnickle and Heikurinen,
15 1989; Boucher et al., 1998; Örländer et al., 1998; Bedford and Sutton, 2000;
16 Thiffault et al., 2004).

17

18 Our results support the assumption that the positive effect of disk
19 scarification might be related, at least in part, to increased soil water availability
20 and uptake by seedlings, that leads to enhanced seedling water relations. The
21 most direct influence of scarification on soil water is through decreased
22 interception and competition, which results in more water available for planted
23 seedlings (Prévost, 1992; Morris et al., 1993; Burgess et al., 1995; Boucher et al.,

1 1998; Brais, 2001). Decreased rainfall interception following scarification can also
2 be the consequence of reduced woody debris and humus thickness (Prévost,
3 1992). Enhanced soil temperature following disk scarification should also
4 contribute to higher seedling water functions and growth, through increased root
5 growth, and root water and nutrient absorption (Bassman, 1989; Bowen, 1991;
6 Lyr and Garbe, 1995; Boucher et al., 1998, 2001). Soil temperature, although not
7 measured in the present study, may be an important growth-limiting factor in
8 intact LW because of the high albedo of the lichen mat (Kershaw and Rouse,
9 1971).

10

11 Other important changes resulting from disk scarification include the
12 diminished influence of competition on nutrient availability, ectomycorrhizal
13 interference on seedling roots, and allelopathic influence (Lanini and Radosevich,
14 1986; Mallik, 1993; Zhu and Mallik, 1994; Bradley et al., 1997; Yamasaki et al.,
15 1998, 2002). Decreased light interception by competition due to scarification
16 normally contributes to seedling response, but this influence was probably
17 moderate in the present study, since light availability was relatively high even in
18 intact LW plots, with 78% of full light available at seedling height (Girard, 2004).
19 Absence of a positive effect by patch scarification on seedling growth might be
20 simply because of the mildness of this manual scarification method. The method
21 used in the study removed the aerial part of competing vegetation in a 15-cm
22 radius only, and decreased partially the humus layer, rarely reaching the mineral
23 soil.

1 Values of stomatal conductance for water vapor in black spruce were
2 essentially unaffected by treatments, contrary to those in jack pine that were
3 strongly reduced in the LWno and the LWps treatments. Black spruce is much
4 more shade tolerant than jack pine, and a lack of physiological plasticity is a
5 general feature of shade tolerant tree species (Givnish, 1988; Canham, 1989;
6 Bazzaz and Wayne, 1994; Walters and Reich, 2000). Black spruce robustness to
7 contrasting soil water conditions has been documented in Zine El Abidine et al.
8 (1995), but Grossnickle and Blake (1986) found that black spruce seedlings are
9 more sensitive than jack pine to atmospheric humidity deficit. However, since
10 atmospheric conditions were comparable among treatments (Girard, 2004), we
11 believe that soil water and possibly soil temperature (Boucher et al., 2001) were
12 the variables controlling the seedling's water relations on our sites. A portion of
13 the higher growth rate observed in jack pine could therefore be attributed to its
14 greater water stress tolerance, expressed as higher values of stomatal
15 conductance and of predawn water potential compared to black spruce
16 (Grossnickle and Blake, 1986; Raison et al., 1992; Bernier, 1993). Furthermore,
17 Tan et al. (1992) and Stewart et al. (1995) suggest that the drought response of
18 black spruce is to maximize the rate of photosynthesis, while that of jack pine
19 may be to optimize photosynthesis in balance with water loss. The former
20 response would be better suited to shorter and less frequent drought events, and
21 the latter more suited to continuous or recurring drought, such as in lichen
22 woodlands (Bergeron 1996).

23

24

1 4.2. Comparison of treated lichen woodlands with managed black spruce-
2 feathermoss stands

3

4 Growth and water relations of the planted seedlings (both species)
5 associated with the disk scarified LW plots were generally comparable to those
6 associated with the managed black spruce-feathermoss stands (BSFM), although
7 significant differences in seedling growth suggest higher productivity in BSFM. In
8 spite of the usual positive impact of disk scarification on seedling growth and
9 survival, nutritional constraints may further limit seedling growth on the scarified
10 LW sites. A parallel study on the same sites has found a lower foliar nitrogen of
11 black spruce seedlings in disk scarified LW compared to BSFM sites. Results
12 from the same study also suggest an influence of the residual plant competition
13 on these nutritional differences (Girard, 2004). The higher relative biomass
14 allocation to roots in seedlings in LW compared to that in BSFM may be an
15 additional indication of greater resource limitations in the soils of LW (Grime,
16 1994; Canham et al., 1996).

17

18

19 4.3. Silvicultural implications

20

21 The present study constitutes a first step at the determination of optimal
22 practices for the afforestation of black spruce lichen woodlands. One important
23 issue in this particular challenge is the planting check caused by water stress,

1 because of the putative – but still not documented – xeric conditions that
2 characterize LW (Burdett et al. 1984; Bergeron, 1996). Our results indicate that
3 soil water availability in intact or lightly prepared patch scarified LW is likely a
4 critical variable for early seedling growth. However, in LW prepared with disk
5 scarification, this constraint appears to be significantly reduced, but nutritional
6 constraints generated by the residual plant competition may still limit growth. The
7 level of disturbance before planting necessary to achieve full seedling growth
8 potential in LW should be further investigated, for example by combining harvest
9 with scarification before planting, or by testing more aggressive site preparation
10 methods (Prévost, 1996; Örlander et al., 1998; Brais, 2001). Our results also
11 suggest that jack pine could constitute a wise silvicultural choice over black
12 spruce, especially with LW established on highly drought-prone sites such as
13 sandy deposits or steep south-facing slopes.

1 **Acknowledgements**

2

3 We thank François Girard, Dominic Dufour, Michaël Tremblay, Geneviève
4 Gobeil and Sébastien Audet for their assistance in the field, and Marie-Claude
5 Lambert for statistical advice on the MIXED procedure. Comments by three
6 anonymous reviewers were of great help in increasing the quality of the paper.
7 Logistical help from Jacques Allaire, Denis Walsh, and Damien Côté was also
8 greatly appreciated, as well as from Hubert Morin and François Gionest for the
9 meteorological data in a nearby station. Research funding was provided by the
10 FQRNT (Technological and Natural Research Quebec funds) and by the
11 *Ministère des Ressources Naturelles du Québec, Région 02.*

12

1 **References**

2

3 Anonymous, 2005. Ministère des Ressources naturelles, de la faune et des
4 parcs. Direction des inventaires forestiers, 3^e inventaire écoforestier.

5

6 Bassman, J.H., 1989. Influence of two site preparation treatments on
7 ecophysiology of planted *Picea engelmannii* X *glauca* seedlings. Can. J.
8 For. Res. 19, 1359-1370.

9

10 Bauce, E., Crépin, M., Carisey, N., 1994. Spruce budworm growth, development
11 and food utilization on young and old balsam fir trees. Oecologia 97, 499-
12 507.

13

14 Bazzaz, F.A., Wayne, P.M., 1994. Coping with environmental heterogeneity: The
15 physiological ecology of tree seedling regeneration across the gap-
16 understory continuum. In: Caldwell, M.M., Pearcy, R.W. (Eds.),
17 Exploitation of Environmental Heterogeneity by Plants. Ecophysiological
18 Processes Above- and Belowground. Academic Press, Inc., pp. 349-390.

19

20 Bedford, L., Sutton, R.F., 2000. Site preparation for establishing lodgepole pine in
21 the sub-boreal spruce zone of interior British Columbia: the Bednesti trial,
22 10-year results. For. Ecol. Manage. 126, 227-238.

23

- 1 Bergeron, J.-F., 1996. Domaine de la pessière noire à mousses. In: Bérard, J.A.,
2 Côté, M. (Eds.), Manuel de Foresterie. Les Presses de l'Université Laval,
3 Québec,, pp. 223-238.
- 4
- 5 Bernier P. Y., 1993. Comparing natural and planted black spruce seedlings. I.
6 Water relations and growth. Can. J. For. Res. 23, 2424-2434.
- 7
- 8 Bernier, P.Y., Raulier, F., Stenberg, P., Ung, C.H., 2001. Importance of needle
9 age and shoot structure on canopy net photosynthesis of balsam fir
10 (*Abies balsamea*): a spatially inexplicit modeling analysis. Tree Physiol.
11 21, 815-830.
- 12 Boucher, J.F., Wetzel, S., Munson, A.D., 1998. Leaf level response of planted
13 eastern white pine (*Pinus strobus* L.) seven years after intensive
14 silvicultural treatments. For. Ecol. Manage. 107, 291-307.
- 15
- 16 Boucher, J.-F., Bernier, P.Y., Munson, A.D., 2001. Radiation and soil
17 temperature interactions on the growth and physiology of eastern white
18 pine (*Pinus strobus* L.) seedlings. Plant and Soil 236, 165-174.
- 19
- 20 Bowen, G.D., 1991. Soil temperature, root growth, and plant function. In: Waisel,
21 Y., Eshel, A., Kafkafi, U. (Eds.), Plant Roots: The Hidden Half. Marcel
22 Dekker, Inc., New-York, pp. 309-330.
- 23

- 1 Bradley, R.L., Fyles, J.W., Titus, B., 1997. Interactions between *Kalmia* humus
2 quality and chronic low C inputs in controlling microbial and soil nutrient
3 dynamics. *Soil Biol. Biochem.* 29, 1275-1283.
- 4
- 5 Brais, S., 2001. Persistence of soil compaction and effects on seedling growth in
6 northwestern Quebec. *Soil Sci. Soc. Am. J.* 65, 1263-1271.
- 7
- 8 Burdett, A.N., Herring, L.J., Thompson, C.F., 1984. Early growth of planted
9 spruce. *Can. J. For. Res.* 14, 644-651.
- 10
- 11 Burgess, D., Baldock, J.A., Wetzel, S., Brand, D.G., 1995. Scarification,
12 fertilization and herbicide treatment effects on planted conifers and soil
13 fertility. *Plant and Soil* 168-169, 513-522.
- 14
- 15 Canham, C.D., 1989. Different responses to gaps among shade-tolerant tree
16 species. *Ecology* 70, 548-550.
- 17
- 18 Canham, C.D., Berkowitz, A.R., Kelly, V.R., Lovett, G.M., Ollinger, S.V., Schnurr,
19 J., 1996. Biomass allocation and multiple resource limitation in tree
20 seedlings. *Can. J. For. Res.* 26, 1521-1530.
- 21
- 22 Devore, J., Peck, R., 1994. *Introductory Statistics*. 2nd edition. St. Paul, West
23 Publishing Company.

- 1
- 2 Environment Canada, 2004. Climatic Normals and Means, Chibougamau-
- 3 Chapais, year 1971-2000. Atmospheric environment service, Environment
- 4 Canada.
- 5
- 6 Fleming, R.L., Black, T.A., Eldridge, N.R., 1994. Effects of site preparation on
- 7 root zone soil water regimes in high-elevation forest clearcuts. For. Ecol.
- 8 Manage. 68, 173-188.
- 9
- 10 Gagnon, R., Morin, H., 2001. Les forêts d'épinette noire du Québec: dynamique,
- 11 perturbations et biodiversité. Le Naturaliste Canadien 125, 26-35.
- 12
- 13 Girard, F. 2004. Interaction Compétitive entre des Plants d'Épinette Noire, des
- 14 Lichens et des Éricacées dans les Pessières à Lichens Trois Ans après
- 15 Traitements de Préparation de Terrain. Master degree thesis. Université
- 16 du Québec à Chicoutimi, Qc, Canada.
- 17
- 18 Givnish, T.J., 1988. Adaptation to sun and shade: A whole-plant perspective.
- 19 Aust. J. Plant Physiol. 15, 63-92.
- 20
- 21 Grime, J.P., 1994. The role of plasticity in exploiting environmental heterogeneity.
- 22 In: Caldwell, M.M., Pearcy, R.W. (Eds.), Exploitation of Environmental
- 23 Heterogeneity by Plants. Ecophysiological Processes Above- and

- 1 Belowground. Academic Press, Inc., pp. 1-19.
- 2
- 3 Grossnickle, S.C., Blake, T.J., 1986. Environmental and physiological control of
- 4 needle conductance for bare-root black spruce, white spruce, and jack
- 5 pine seedlings on boreal cutover sites. Can. J. Bot. 64, 943-949.
- 6
- 7 Grossnickle, S.C., Heikurinen, J., 1989. Site preparation: water relations and
- 8 growth of newly planted jack pine and white spruce. New For. 3, 99-123.
- 9
- 10 Harvey, B., Brais, S., 2002. Effects of mechanized careful logging on natural
- 11 regeneration and vegetation competition in the southeastern Canadian
- 12 boreal forest. Can. J. For. Res. 32, 653-666.
- 13
- 14 Inderjit, Mallik, A.U., 1996. Growth and physiological responses of black spruce
- 15 (*Picea mariana*) to sites dominated by *Ledum groenlandicum*. J. Chem.
- 16 Ecol. 22, 575-585.
- 17
- 18 Inderjit, Mallik, A.U., 2002. Can *Kalmia angustifolia* interference to black spruce
- 19 (*Picea mariana*) be explained by allelopathy? For. Ecol. Manage. 160, 75-
- 20 84.
- 21
- 22 Kershaw, K.A., Rouse, W.R., 1971. Studies on lichen-dominated systems. II The
- 23 growth pattern of *Cladonia alpestris* and *Cladonia rangiferina*. Can. J. Bot.

- 1 49, 1401-1410.
- 2
- 3 Lanini W.T., Radosevich S.R., 1986. Response of three conifer species to site
4 preparation and shrub control. For. Sci. 32, 61-77.
- 5
- 6 Lyr H., V. Garbe, 1995. Influence of root temperature on growth of *Pinus*
7 *sylverstris*, *Fagus sylvatica*, *Tilia cordata* and *Quercus robur*. Trees 9, 210-
8 213.
- 9
- 10 Mäkitalo, K., Hyvönen, J., 2004. Late summer soil water content on clear-cut
11 reforestation areas two decades after site preparation in Finnish Lapland.
12 For. Ecol. Manage. 189, 57-75.
- 13
- 14 Mallik, A.U., 1991. Cutting, burning, and mulching to control *Kalmia*: results of a
15 greenhouse experiment. Can. J. For. Res. 21, 417-420.
- 16
- 17 Mallik, A.U., 1993. Ecology of a forest weed of Newfoundland: vegetative
18 regeneration strategy of *Kalmia angustifolia*. Can. J. Bot. 71, 161-166.
- 19
- 20 Morris, L.A., Moss, S.A., Garbett, W.S., 1993. Competitive interference between
21 selected herbaceous and woody plants and *Pinus taeda* L during 2
22 growing seasons following planting. For. Sci. 39, 166-187.
- 23

- 1 Nelson, D.W., Sommers, L.E., 1983. Total carbon, organic carbon and organic
2 matter. In: Page, A.L., Miller, R.H., Keeney, D.R. (Eds.), Methods of Soil
3 Analysis, Part 3-Chemical Methods, second ed. American society of
4 agronomy and soil science society of America, Wisconsin, pp. 961-1001.
5
- 6 Örlander, G., 1986. Effect of planting and scarification on the water relations in
7 planted seedlings of Scots pine. Studia Fores. Suec. 173, 1-17.
8
- 9 Örlander, G., Hallsby, G., Gemmel, P., Wilhelmsson, C., 1998. Inverting
10 improves establishment of *Pinus contorta* and *Picea abies*-10-year results
11 from a site preparation trial in Northern Sweden. Scand. J. For. Res. 13,
12 160-168.
- 13
- 14 Parker, W.C., Watson, S.R., Cairns, D.W., 1997. The role of hair-cap mosses
15 (*Polytrichum* spp.) in natural regeneration of white spruce (*Picea glauca*
16 (Moench) Voss). For. Ecol. Manage. 92, 19-28.
- 17
- 18 Payette S., 1992. Fire as a controling prosscess in the north American boreal
19 forest. In: Shugart, H.H., Leemans, R., Bonan, G.B. (Eds.), Systems
20 Analysis of the Global Boreal Forest. Cambridge, pp. 144-169.
- 21
- 22 Payette, S., Bhiry, N., Delwaide, A., Simard, M., 2000. Origin of the lichen
23 woodland at its southern range limit in eastern Canada: the catastrophic

- 1 impact of insect defoliators and fire on the spruce-moss forest. Can. J. For.
2 Res. 30, 288-305.
- 3
- 4 Poorter H., Lewis. C., 1986. Testing differences in relative growth rate: A method
5 avoiding curve fitting and pairing. Physiol. Plant. 67, 223-226.
- 6
- 7 Prévost, M., 1992. Effets du scarifiage sur les propriétés du sol, la croissance
8 des semis et la compétition: revue des connaissances actuelles et
9 perspectives de recherches au Québec. Ann. Sci. For. 49, 277-296.
- 10
- 11 Prévost M., 1996. Effet du scarifiage sur les propriétés du sol et
12 l'ensemencement naturel dans une pessière noire à mousses de la forêt
13 boréale québécoise. Can. J. For. Res. 26, 72-86.
- 14
- 15 Raison, R.J., Khanna, P.K., Benson, M.L., Myers, B.J., McMurtrie, R.E., Lang,
16 A.R.G., 1992. Dynamics of *Pinus radiata* foliage in relation to water and
17 nitrogen stress. II. Needle loss and temporal changes in total foliage mass.
18 For. Ecol. Manage. 52, 159-178.
- 19
- 20 Riverin, S., Gagnon, R., 1996. Dynamique de la régénération d'une pessière à
21 lichen dans la zone de la pessière noire à mousse, nord du Saguenay-
22 Lac-Saint-Jean (Québec). Can. J. For. Res. 26, 1504-1509.
- 23

- 1 Rundel, P.W., Jarrell W.M., 1991. Water in the environment. In: Pearcy R.W.,
2 Ehleringer J.R., Mooney H.A., Rundel P.W. (Eds.), Plant Physiological
3 Ecology: Field Methods and Instrumentation. London, pp. 15-29.
- 4
- 5 Saucier, J.P., Berger, J.P., D'Avignon, H., Racine, P., 1994. Le Point
6 d'Observation Écologique. Ministère des Ressources Naturelles du
7 Québec, Québec.
- 8
- 9 Scholander P.F., Hammel H.T., Bradstreet E.D., Hemmingsen E.A., 1965. Sap
10 pressure in vascular plants. Sci. 148, 339-346.
- 11
- 12 Sirois, L., 1993. Impact of fire on *Picea mariana* and *Pinus banksiana* seedlings
13 in subarctic lichen woodlands. J. Veg. Sci. 4, 795-802.
- 14
- 15 Stathers, R.J., Spittlehouse, D.L., 1990. Forest soil temperature manual. B.C.
16 Min. For., Victoria B.C., FRDA report 130.
- 17
- 18 Steel, R.G.D., Torrie, J.H., 1980. Principles and Procedures of Statistics: a
19 Biometrical Approach, second ed. McGraw-Hill Publishing Company, New-
20 York.
- 21
- 22 Steijlen, I., Nilsson, M.-C., Zackrisson, O., 1995. Seed regeneration of scots pine
23 in boreal forest stands dominated by lichen and feather moss. Can. J. For.

- 1 Res. 25, 713-723.
- 2
- 3 Stewart, J.D., Abidine, A.Z.E., Bernier, P.Y., 1995. Stomatal and mesophyll
4 limitations of photosynthesis in black spruce seedlings during multiple
5 cycles of drought. *Tree Physiol.* 15, 57-64.
- 6
- 7 Tan, W.X., Blake, T.J., Boyle, T.J.B., 1992. Drought Tolerance in Faster-Growing
8 and Slower-Growing Black Spruce (*Picea mariana*) Progenies .1. Stomatal
9 and Gas Exchange Responses to Osmotic Stress. *Physiol. Plant.* 85, 639-
10 644.
- 11
- 12 Thiffault, N., Jobidon R., Munson, A.D., 2003. Performance and physiology of
13 large containerized and bare-root spruce seedlings in relation to
14 scarification and competition in Québec (Canada). *Ann. For. Sci.* 60, 645-
15 655.
- 16
- 17 Thiffault, N., Cyr, G., Prégent, G., Jobidon, R., Charrette, L., 2004. Régénération
18 artificielle des pessières noires à éricacées : effets du scarifiage, de la
19 fertilisation et du type de plants après 10 ans. *For. Chron.* 80, 141-149.
- 20
- 21 Tremblay, P.S., 1996. Opérations forestières. In: Bérard, J.A., Côté, M. (Eds.),
22 Manuel de Foresterie. Les Presses de l'Université Laval, Québec, pp.
23 1177-1225.

- 1 Walters, M.B., Reich, P.B., 2000. Trade-offs in low-light CO₂ exchange: a
2 component of variation in shade tolerance among cold temperate tree
3 seedlings. Func. Ecol. 14, 155-165.
- 4
- 5 Yamasaki S. H., Fyles J. W., Egger K. N., Titus B. D., 1998. The effect of *Kalmia*
6 *angustifolia* on the growth, nutrition and ectomycorrhiza symbiont
7 community of black spruce. For. Ecol. Manage. 105, 197-207.
- 8
- 9 Yamasaki, S.H., Fyles, J.M., Titus, B.D., 2002. Interaction among *Kalmia*
10 *angustifolia*, soil characteristics, and the growth and nutrition of black
11 spruce seedlings in two boreal Newfoundland plantations of contrasting
12 fertility. Can. J. For. Res. 32, 2215-2224.
- 13
- 14 Zar, J.H., 1999. Biostatistical Analysis, fourth ed. Prentice Hall, Englewood Cliffs,
15 New Jersey.
- 16
- 17 Zhu, H., Mallik, A.U., 1994. Interactions between *Kalmia* and black spruce:
18 Isolation and identification of allelopathic compounds. J. Chem. Ecol. 20,
19 407-421.
- 20
- 21 Zine El Abidine, A., Stewart, J.D., Bernier, P.Y., Plamondon, A.P., 1995. Diurnal
22 and seasonal variations in gas exchange and water relations of lowland
23 and upland black spruce ecotypes. Can. J. Bot. 73, 716-72.

